

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Box Patent Application
Assistant Commissioner For Patents
Washington, D.C. 20231

NEW APPLICATION TRANSMITTAL

Transmitted herewith for filing is the patent application of Linda B. Couto, Peter C. Colosi and Xiaobing Qian for **ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS**.

CERTIFICATION UNDER 37 C.F.R. § 1.10

I hereby certify that this New Application Transmittal and the documents referred to as enclosed therein are being deposited with the U.S. Postal Service on this date **December 22, 1999** in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number **EL 079 496 941 US** addressed to: **Box Patent Application**, Assistant Commissioner For Patents, Washington, D.C. 20231.

Marlene Garitano
Marlene Garitano

1. **Type Of Application**

This new application is for a(n)

- ☒ Original (nonprovisional)
☒ Continuation-In-Part (C-I-P).

2. **Benefit Of Prior U.S. Application(s)** (35 U.S.C. §§ 119(e), 120, or 121)

- ☒ The new application being transmitted claims the benefit of prior U.S. application(s) and enclosed are **ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR US APPLICATION(S) CLAIMED**.

3. **Papers Enclosed That Are Required For Filing Date Under 37 C.F.R. § 1.53(b) (Regular) or 37 C.F.R. § 1.153 (Design) Application**

- 64 Pages of Specification
6 Pages of Claims
1 Page of Abstract
15 Sheets of Informal Drawings

4. **Additional Papers Enclosed**

- ☒ Submission of "Sequence Listing," computer readable copy and/or amendment pertaining thereto for biotechnology invention containing nucleotide and/or amino acid sequence.
☒ Certificate Re: Sequence Listing

5. **Declaration**

- ☒ Enclosed
☒ Unexecuted

6. **Inventorship Statement**

The inventorship for all the claims in this application is:

- ☒ the same

7. **Language**

- ☒ English

8. **Assignment**

- ☒ An assignment of the invention to **Avigen, Inc.** is attached
☒ Unexecuted.
☒ Form PTO-1595 will follow.

jc525 U.S. PTO
09/470618
12/22/99

9. Fee Calculation (37 C.F.R. § 1.16)

☒ Regular application

CLAIMS AS FILED

Number Filed	Number Extra	Rate	Basic Fee - \$760.00 (37 C.F.R. § 1.16(a))
Total Claims (37 C.F.R. § 1.16(c))	40 - 20 =	20 × \$18.00 =	\$360.00
Independent Claims (37 C.F.R. § 1.16(b))	4 - 3 =	1 × \$78.00 =	\$78.00
Multiple Dependent Claim(s), if any (37 C.F.R. § 1.16(d))	+ \$260.00 =		\$0.00
Filing Fee Calculation			\$1198.00

10. Small Entity Statement(s)

- ☒ Verified Statement(s) that this is a filing by a small entity under 37 C.F.R. §§ 1.9 and 1.27 is(are) attached.
- ☒ Unexecuted.

Filing Fee Calculation (50% of above) \$599.00

11. Fee Payment Being Made At This Time

- ☒ Enclosed
- ☒ basic filing fee

\$599.00

Total Fees Enclosed \$599.00

12. Method of Payment of Fees

- ☒ Check in the amount of \$599.00

13. Authorization To Charge Additional Fees and Credit Overpayment

- ☒ The Commissioner is hereby authorized to charge payment of any fees associated with this communication or credit any overpayment to Deposit Account No. 08-1290. An originally executed duplicate of this transmittal is enclosed for this purpose


14. Power of Attorney by Assignee

- ☒ Enclosed
- ☒ Unexecuted.

15. Return Receipt Postcard

- ☒ Enclosed

Dated: December 22, 1999


Kamrin T. MacKnight
Registration No. 38,230

MEDLEN & CARROLL, LLP
220 Montgomery Street, Suite 2200
San Francisco, California 94104
(415) 705-8410

☒ Incorporation By Reference Of Added Pages

- ☒ Plus Added Pages For New Application Transmittal Where Benefit Of Prior U.S. Application(s) Claimed

Number of pages added 1

ADDED PAGES FOR APPLICATION TRANSMITTAL
WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED

16. Relate Back

A. 35 U.S.C. § 119(e)

B. 35 U.S.C. §§ 120, 121 and 365(c)

- ☒ Amend the Specification by inserting before the first line the sentence: "This application is a Continuation-in-Part of U.S. Patent Application Ser. No. 09/364,862, filed July 30, 1999, which claims benefit of U.S. provisional application Nos 60/125,974 filed March 24, 1999 and 60/104,994 filed October 20, 1998, all of which are hereby incorporated by reference in their entireties."

Applicant / Patentee: Linda B. Couto *et al.*

For: **ADENO-ASSOCIATED VECTORS FOR
EXPRESSION OF FACTOR VIII BY TARGET CELLS**

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR § 1.9(c) - SMALL BUSINESS CONCERN)**

I hereby declare that I am an official of the small business concern empowered to act on behalf of the concern identified below:

Avigen, Inc.
1201 Harbor Bay Parkway, #1000
Alameda, California 94502

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR §§ 121.3-18, and reproduced in 37 CFR § 1.9(d), for purposes of paying reduced fees under §§ 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention entitled **ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS** by inventors named **Linda B. Couto *et al.***, described in the specification filed herewith.

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR § 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Date: _____

By: _____

Name: _____

Title: _____

ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS

This application is a Continuation-in-Part of U.S. Patent Application Ser. No. 09/364,862, filed July 30, 1999, which claims benefit of U.S. provisional application Nos. 60/125,974 filed March 24, 1999 and 60/104,994 filed October 20, 1998, all of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia gene therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids encoding Factor VIII into a recipient subject suffering from hemophilia A, such that the subject's blood is able to clot.

BACKGROUND

Hemophilia is a genetic disease characterized by a blood clotting deficiency. In hemophilia A (classic hemophilia, Factor VIII deficiency), an X-chromosome-linked genetic defect disrupts the gene encoding Factor VIII, a plasma glycoprotein, which is a key component in the blood clotting cascade. Human Factor VIII is synthesized as a single chain polypeptide, with a predicted molecular weight of 265 kDa. The Factor VIII gene codes for 2351 amino acids, and the protein has six domains, designated from the amino to the carboxy terminus as A1-A2-B-A3-C1-C2 (Wood *et al.*, Nature 312:330 [1984]; Vehar *et al.*, Nature 312:337 [1984]; and Toole *et al.*, Nature 312:342 [1984]). Human Factor VIII is processed within the cell to yield a heterodimer primarily comprised of a heavy chain of 200 kDa containing the A1, A2, and B domains and an 80 kDa light chain containing the A3, C1, and C2 domains (Kaufman *et al.*, J. Biol. Chem., 263:6352-6362 [1988]). Both the single chain polypeptide and the heterodimer circulate in the plasma as inactive precursors (Ganz *et al.*, Eur. J. Biochem., 170:521-528 [1988]). Activation of Factor VIII in plasma is initiated by thrombin cleavage between the A2 and B domains, which releases the B domain and

results in a heavy chain consisting of the A1 and A2 domains. The 980 amino acid B domain is deleted in the activated procoagulant form of the protein. Additionally, in the native protein, two polypeptide chains ("a" and "b"), flanking the B domain, are bound to a divalent calcium cation. Hemophilia may result from point mutations, deletions, or mutations resulting in a stop codon (*See, Antonarakis et al., Mol. Biol. Med., 4:81 [1987]*).

The disease is relatively rare, afflicting approximately one in 10,000 males. Hemophilia in females is extremely rare, although it may occur in female children of an affected father and carrier mother, as well as in females with X-chromosomal abnormalities (*e.g., Turner syndrome, X mosaicism, etc.*). The severity of each patient's disease is broadly characterized into three groups--"mild," "moderate," and "severe," depending on the severity of the patient's symptoms and circulating Factor VIII levels. While normal levels of Factor VIII range between 50 and 200 ng/mL plasma, mildly affected patients have 6-60% of this value, and moderately affected patients have 1-5% of this value. Severely affected hemophiliacs have less than 1% of normal Factor VIII levels.

While hemophiliacs clearly require clotting factor after surgery or severe trauma, on a daily basis, spontaneous internal bleeding is a greater concern. Hemophiliacs experience spontaneous hemorrhages from early infancy, as well as frequent spontaneous hemarthroses and other hemorrhages requiring clotting factor replacement. Without effective treatment, chronic hemophilic arthropathy occurs by young adulthood. Severely affected patients are prone to serious hemorrhages that may dissect through tissue planes, ultimately resulting in death due to compromised vital organs.

Hematomas are commonly observed in moderately and severely affected hemophiliacs. In these patients, hematomas have a tendency to progressively enlarge and dissect in all directions. Some of these hematomas expand locally, resulting in local compression of adjacent organs, blood vessels, and nerves. A rare, yet often fatal, complication of abdominal hematomas is the perforation and drainage of the hematoma into the colon, resulting in infection and septicemia. Intracranial and/or

extracranial hemorrhage also represent very dangerous bleeding situations. While subcutaneous hematomas may dissect into muscle, pharyngeal and retropharyngeal hematomas (*e.g.*, complicating bacterial or viral pharyngitis) may enlarge and obstruct the airway, sometimes resulting in a life-threatening situation that requires
5 administration of a sufficient dose of Factor VIII concentrate to normalize the Factor VIII level.

In addition to hematomas, hemarthroses are commonly observed in hemophiliacs, with bleeding into the joint accounting for approximately 75% of hemophilic bleeding. Repeated hemorrhaging into the joints eventually results in
10 extensive destruction of articular cartilage, synovial hyperplasia, and other reactive changes in adjacent tissues and bone. A major complication of repeated hemarthroses is joint deformity, which is often accompanied by muscle atrophy and soft tissue contractures; osteoporosis and cystic areas in the subchondral bone may also develop, along with progressive loss of joint space.

Other symptoms are often observed in hemophiliacs, including hematuria and mucous membrane bleeding. Hematuria is experienced by virtually all severely affected hemophiliacs sometime during their lifetimes, and mucous membrane bleeding is common in hemophiliacs. Bone cysts (pseudotumors) are rare, but dangerous complications of hemophilic bleeding. In many of these cases, immediate treatment is
15 necessary.

In the early 1980s, many severely affected hemophiliacs were treated with Factor VIII concentrate about three times weekly. Unfortunately, these concentrates transmitted viruses, such as hepatitis B and/or C, and human immunodeficiency virus (HIV). In the United States and Western Europe, at least 75% of Factor VIII
25 concentrate recipients have been reported to have anti-HIV antibodies (*See*, Schrier and Leung, *supra*). Some of these patients also developed HIV-associated immune thrombocytopenia, a very serious complication in hemophiliacs. In spite of antiviral therapy (*e.g.*, with zidovudine and pentamidine prophylaxis), which has tended to slow disease progression, full-blown AIDS (acquired immunodeficiency syndrome) occurs at
30 an inexorable rate in hemophiliacs infected with HIV. Indeed, this has reversed the

improvement in the life expectancy of hemophiliacs, which peaked at 66 years of age during the 1970s, and has dropped to 49 years (*See*, Schrier and Leung, *supra*). The development of virus-free preparations and recombinant Factor VIII has helped control infectious viral contamination.

5 However, for hemophiliacs, the availability of viral-free concentrates and recombinant Factor VIII, while significant, is but part of the solution. In order to prevent spontaneous internal bleeding episodes, patients suffering from hemophilia A must consistently have serum Factor VIII levels of about 1%, and preferably 5%. Currently, the cost of viral-free concentrates and recombinant Factor VIII make it
10 prohibitively expensive to administer the clotting factor prophylactically or on a maintenance basis. Indeed, most hemophiliacs in the U.S. do not receive recombinant Factor VIII therapy on a maintenance basis, but only receive it prior to activities or events which might cause bleeding (*e.g.*, surgery), or as a treatment for spontaneous bleeding.

15 Moreover, even if cost effective preparations of recombinant or virus-free Factor VIII were available, a steady state level of Factor VIII cannot be achieved by its daily administration. At best, patients receive widely varying levels of Factor VIII. Immediately following the administration, the levels are super-physiological, while prior to administration the levels are sub-physiological. Thus, there remains a need for
20 methods and compositions that are relatively economic, yet effective in the treatment and prevention of bleeding in hemophiliacs, particularly spontaneous bleeds. Furthermore, there is a need in the art for methods and compositions for long term delivery of clotting factors (*e.g.*, Factor VIII) which more closely mimic the steady state physiological levels observed in normal individuals.

SUMMARY OF THE INVENTION

25 The present invention provides improved viral vectors suitable for gene therapy to treat hemophilia. In particular, the present invention provides AAV vectors and methods for treating hemophilia A by delivering nucleic acids coding for the clotting

protein Factor VIII. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the vectors.

The present invention is particularly suited for use in hemophilia A gene therapy. Accordingly, in one embodiment of the invention, at least one AAV vector containing a nucleic acid molecule encoding Factor VIII is operably linked to control sequences that direct expression of Factor VIII in a suitable recipient cell. The AAV vectors are then introduced into a recipient cell of the subject, under conditions that result in expression of Factor VIII. The subject, therefore, has a continuous supply of Factor VIII available to clot blood during bleeding episodes.

Using the methods of the present invention, long term expression of therapeutic levels of Factor VIII have been achieved *in vivo*. In one embodiment, animals were administered, via the portal vein, two AAV vectors: one carrying the DNA sequence coding for the heavy chain of Factor VIII and the other carrying the DNA sequence coding for the light chain of Factor VIII. Blood samples were collected periodically and assayed for Factor VIII activity. Reproducibly, animals expressed between 600 and 900 ng/ml of biologically active Factor VIII, levels that are well above the normal physiological levels of approximately 200 ng/ml. Furthermore, these levels have been sustained for over 13 months without a decrease in Factor VIII levels or activity. In a related embodiment, a B-domain deleted form of Factor VIII was cloned into a single AAV vector and shown to express biologically active Factor VIII.

It is not intended, however, that the present invention be limited to specific embodiments. Many different forms of recombinant Factor VIII have been made and tested both *in vitro* and *in vivo*, using a variety of different control and regulatory sequences. Any DNA sequence coding for biologically active Factor VIII can be expressed using the AAV vectors and methods taught in the present invention. Therefore, the present invention encompasses any AAV vector or vectors containing Factor VIII sequences that produce biologically active Factor VIII protein *in vitro* or *in vivo*.

For example, in some embodiments, the AAV vector contains the first 57 base pairs of the Factor VIII heavy chain which encodes the 10 amino acid signal sequence,

as well as the human growth hormone (hGH) polyadenylation sequence. In some alternative embodiments, the vector also contains the A1 and A2 domains, as well as 5 amino acids from the N-terminus of the B domain, and/or 85 amino acids of the C-terminus of the B domain, as well as the A3, C1, and C2 domains. In yet other
5 embodiments, the nucleic acids coding for Factor VIII heavy chain and light chain were cloned into a single vector separated by 42 nucleic acids coding for 14 amino acids of the B domain.

The present invention also provides methods for administering the above-described vectors. For example, it is intended that the present invention encompass
10 methods suitable for delivery of the AAV vectors to the livers of recipient patients or test animals. It is not intended that the present invention be limited to any particular route of administration. However, in preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal or arterial vasculature.

These and other embodiments of the invention will readily occur to those of
15 ordinary skill in the art in view of the disclosure herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a schematic representation of the Factor VIII protein.

Figure 2 provides a schematic representation of a B-domain deleted form of
20 Factor VIII protein.

Figure 3 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (AAV-F8-1) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 4 provides a schematic representation of a B-domain deleted Factor VIII
25 AAV construct (PVM4.1c-F8AB) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 5 provides the sequence of pAAV-F8-1 (ITR to ITR), with the plasmid backbone omitted.

Figure 6 provides the sequence of pVm4.1cF8ΔB (ITR to ITR), with the
30 plasmid backbone omitted.

Figure 7 provides a map of rAAV-hFVIII-HC and rAAV-hFVIII-LC vectors.

Figure 8 provides a graph demonstrating the expression of various human FVIII constructs in mouse plasma.

Figure 9 provides Southern blot analyses of liver DNA using probes specific
5 for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

Figure 10 provides Southern blot analyses of DNA from different tissues using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

Figure 11 provides Northern blot analyses of liver RNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

10

GENERAL DESCRIPTION OF THE INVENTION

The present invention relates to improved viral vectors useful for expressing gene products at high levels in human cells. In particular, the present invention provides AAV vectors suitable for gene therapy. These vectors are capable of
15 delivering nucleic acid containing constructs which result in the production of Factor VIII protein in a host. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

The AAV vectors and rAAV virions of the present invention can be produced
20 using standard methodology known to those of skill in the art. The methods generally involve the steps of: (1) introducing an AAV vector into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses
25 and/or accessory function vectors into the host cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions. The AAV vector, AAV helper construct and the helper virus or accessory function vector(s) can be introduced into the host cell either
30 simultaneously or serially, using standard transfection techniques.

Unless otherwise indicated, the practice of the present invention employs conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, including those described in such references as Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*; Glover (ed.) *DNA Cloning: A Practical Approach*, Vols. I and II; Gait (ed.) *Oligonucleotide Synthesis*; Hames and Higgins (eds.) *Nucleic Acid Hybridization*; Hames and Higgins (eds.) *Transcription and Translation*; Tijessen (ed.) *CRC Handbook of Parvoviruses*, Vols. I and II; and Fields and Knipe (eds.) *Fundamental Virology*, 2nd Edition, Vols. I and II.

Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

As used herein, the terms "gene transfer" and "gene delivery" refer to methods or systems for reliably inserting a particular nucleotide sequence (*e.g.*, DNA) into targeted cells. In particularly preferred embodiments, the nucleotide sequence comprises at least a portion of Factor VIII.

As used herein, the terms "vector," and "gene transfer vector" refer to any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus, virion, etc., which is capable of replication when associated with the proper control sequences and/or which can transfer nucleic acid sequences between cells. Thus, the term includes cloning and expression vectors, as well as viral vectors.

Gene transfer vectors may include transcription sequences such as polyadenylation sites, selectable markers or reporter genes, enhancer sequences, and other control sequences which allow for the induction of transcription. Such control sequences are described more fully below.

The term "expression vector" as used herein refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular

host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, as well as other sequences. Eukaryotic cells are generally known to utilize promoters (constitutive, inducible or tissue specific), enhancers, and termination and polyadenylation signals, although some elements may be deleted and other elements added without sacrificing the necessary expression.

As used herein, the terms "host" and "expression host" refer to organisms and/or cells which harbor an exogenous DNA sequence (*e.g.*, via transfection), an expression vector or vehicle, as well as organisms and/or cells that are suitable for use in expressing a recombinant gene or protein. It is not intended that the present invention be limited to any particular type of cell or organism. Indeed, it is contemplated that any suitable organism and/or cell will find use in the present invention as a host.

As used herein, the terms "viral replicons" and "viral origins of replication" refer to viral DNA sequences that allow for the extrachromosomal replication of a vector in a host cell expressing the appropriate replication factors. In some embodiments, vectors which contain either the SV40 or polyoma virus origin of replication replicate to high copy number, while vectors which contain the replicons from bovine papillomavirus or Epstein-Barr virus replicate extrachromosomally at low copy number may be utilized in other embodiments.

As used herein, the term "AAV vector" refers to a vector having functional or partly functional ITR sequences. The ITR sequences may be derived from an adeno-associated virus serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. The ITRs, however, need not be the wild-type nucleotide sequences, and may be altered (*e.g.*, by the insertion, deletion or substitution of nucleotides), so long as the sequences retain function provide for functional rescue, replication and packaging. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or part, preferably the *rep* and/or *cap* genes but retain functional flanking ITR sequences. Functional ITR sequences are necessary for the rescue, replication and packaging of the AAV virion. Thus, an "AAV vector" is

defined herein to include at least those sequences required in *cis* for replication and packaging (e.g., functional ITRs) of the virus.

As used herein, the term "ITR" refers to inverted terminal repeats. The terms "adeno-associated virus inverted terminal repeats" or "AAV ITRs" refer to the art-recognized palindromic regions found at each end of the AAV genome which function together in *cis* as origins of DNA replication and as packaging signals for the virus. For use in some embodiments of the present invention, flanking AAV ITRs are positioned 5' and 3' of one or more selected heterologous nucleotide sequences. Optionally, the ITRs together with the *rep* coding region or the Rep expression product provide for the integration of the selected sequences into the genome of a target cell.

As used herein, the term "AAV *rep* coding region" refers to the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. Muzyczka (Muzyczka, Curr. Top. Microbiol. Immunol., 158:97-129 [1992]) and Kotin (Kotin, Hum. Gene Ther., 5:793-801 [1994]) provide additional descriptions of the AAV *rep* coding region, as well as the *cap* coding region described below. Suitable homologues of the AAV *rep* coding region include the human herpesvirus 6 (HHV-6) *rep* gene which is also known to mediate AAV-2 DNA replication (Thomson *et al.*, Virol., 204:304-311 [1994]).

As used herein, the term "AAV *cap* coding region" refers to the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These *cap* expression products supply the packaging functions which are collectively required for packaging the viral genome.

As used herein, the term "AAV helper function" refers to AAV coding regions capable of being expressed in the host cell to complement AAV viral functions missing from the AAV vector. Typically, the AAV helper functions include the AAV

rep coding region and the AAV *cap* coding region. An "AAV helper construct" is a vector containing AAV coding regions required to complement AAV viral functions missing from the AAV vector (*e.g.*, the AAV *rep* coding region and the AAV *cap* coding region).

5 As used herein, the terms "accessory functions" and "accessory factors" refer to functions and factors that are required by AAV for replication, but are not provided by the AAV virion (or rAAV virion) itself. Thus, these accessory functions and factors must be provided by the host cell, a virus (*e.g.*, adenovirus or herpes simplex virus), or another expression vector that is co-expressed in the same cell. Generally, the E1,
10 E2A, E4 and VA coding regions of adenovirus are used to supply the necessary accessory function required for AAV replication and packaging (Matsushita *et al.*, Gene Therapy 5:938 [1998]).

As used herein, the term "wild type" ("wt") refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a
15 naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that
20 naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "AAV virion" refers to a complete virus particle, such as a "wild-type" (wt) AAV virus particle (comprising a linear, single-stranded AAV nucleic acid genome associated with an AAV capsid protein coat). In this regard,
25 single-stranded AAV nucleic acid molecules of either complementary sense (*e.g.*, "sense" or "antisense" strands), can be packaged into any one AAV virion and both strands are equally infectious.

As used herein, the terms "recombinant AAV virion," and "rAAV virion" refer to an infectious viral particle containing a heterologous DNA molecule of interest (*e.g.*,

Factor VIII sequence) which is flanked on both sides by AAV ITRs. In some embodiments of the present invention, an rAAV virion is produced in a suitable host cell which contains an AAV vector, AAV helper functions and accessory functions introduced therein. In this manner, the host cell is rendered capable of encoding AAV polypeptides that are required for packaging the AAV vector containing a recombinant nucleotide sequence of interest, such as at least a portion of Factor VIII or portions of Factor VIII domains, into recombinant virion particles for subsequent gene delivery.

As used herein, the term "transfection" refers to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art (*See e.g.*, Graham *et al.*, *Virol.*, 52:456 [1973]; Sambrook *et al.*, *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York [1989]; Davis *et al.*, *Basic Methods in Molecular Biology*, Elsevier, [1986]; and Chu *et al.*, *Gene* 13:197 [1981]. Such techniques can be used to introduce one or more exogenous DNA moieties, such as a gene transfer vector and other nucleic acid molecules, into suitable recipient cells.

As used herein, the terms "stable transfection" and "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA.

As used herein, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

As used herein, the term "transduction" denotes the delivery of a DNA molecule to a recipient cell either *in vivo* or *in vitro*, via a replication-defective viral vector, such as via a recombinant AAV virion.

As used herein, the term "recipient cell" refers to a cell which has been transfected or transduced, or is capable of being transfected or transduced, by a nucleic acid construct or vector bearing a selected nucleotide sequence of interest (*i.e.*, Factor VIII). The term includes the progeny of the parent cell, whether or not the progeny are identical in morphology or in genetic make-up to the original parent, so long as the selected nucleotide sequence is present.

The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, synthetic sequences having codons different from the native gene). Similarly, a cell transfected with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to heterologous DNA, as used herein.

As used herein, "coding sequence" or a sequence which "encodes" a particular antigen, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo*, when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

As used herein, the term "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues

of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

As used herein, the term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

As used herein, the term "regulatory element" refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

The term DNA "control sequences" refers collectively to regulatory elements such as promoter sequences, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate recipient cell.

Transcriptional control signals in eukaryotes generally comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (Maniatis *et al.*, Science 236:1237 [1987]). Promoter and enhancer elements have been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control sequences, *i.e.*, promoters, are also found in prokaryotes). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest (*i.e.*, Factor VIII). Some eukaryotic promoters and enhancers have a broad host range while others are functional in a limited subset of cell types (*See e.g.*, Voss *et al.*, Trends Biochem. Sci., 11:287 [1986]; and Maniatis *et al.*, *supra*, for reviews). For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (Dijkema *et al.*, EMBO J. 4:761 [1985]). Two other examples of promoter and enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 α gene (Uetsuki *et al.*, J. Biol. Chem., 264:5791 [1989]; Kim *et al.*, *Gene* 91:217 [1990]; and Mizushima and Nagata, Nucl. Acids. Res., 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777 [1982]) and the human cytomegalovirus (Boshart *et al.*, Cell 41:521 [1985]). Promoters and enhancers can be found naturally alone or together. For example, the long terminal repeats of retroviruses contain both promoter and enhancer functions. Moreover, generally promoters and enhancers act independently of the gene being transcribed or translated. Thus, the enhancer and promoter may be "endogenous" or "exogenous" or "heterologous." An "endogenous" enhancer/promoter is one which is naturally linked with a given gene in the genome. An "exogenous" or "heterologous" enhancer and promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter.

As used herein, the term "tissue specific" refers to regulatory elements or control sequences, such as a promoter, enhancers, etc., wherein the expression of the nucleic acid sequence is substantially greater in a specific cell type(s) or tissue(s). In particularly preferred embodiments, the albumin promoter and the transthyretin promoter display increased expression of FVIII in hepatocytes, as compared to other cell types. It is not intended, however, that the present invention be limited to the albumin or transthyretin promoters or to hepatic-specific expression, as other tissue specific regulatory elements, or regulatory elements that display altered gene expression patterns, are contemplated.

The presence of "splicing signals" on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells requires expression of signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site" or "poly A sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript.

Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A signal is contained on a 237 bp *Bam*HI/*Bcl*II restriction fragment and directs both termination and polyadenylation (Sambrook *et al.*, *supra*, at 16.6-16.7).

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids are nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. The isolated nucleic acid, oligonucleotide, or polynucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid, oligonucleotide or polynucleotide is to be utilized to express a protein, the oligonucleotide or polynucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide or polynucleotide may single-stranded), but may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide or polynucleotide may be double-stranded).

As used herein, the term "purified" or "to purify" refers to the removal of contaminants from a sample. For example, antibodies may be purified by removal of contaminating non-immunoglobulin proteins; they may also purified by the removal of immunoglobulin that does not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII). The removal of non-immunoglobulin proteins and/or the removal of

immunoglobulins that do not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII) results in an increase in the percent of desired antigen-reactive immunoglobulins in the sample. In another example, recombinant polypeptides of Factor VIII are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant polypeptides is thereby increased in the sample.

As used herein, the term "chimeric protein" refers to two or more coding sequences obtained from different genes, that have been cloned together and that, after translation, act as a single polypeptide sequence. Chimeric proteins are also referred to as "hybrid proteins." As used herein, the term "chimeric protein" refers to coding sequences that are obtained from different species of organisms, as well as coding sequences that are obtained from the same species of organisms.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The composition may comprise an aqueous solution.

As used herein, the term "at risk" is used in references to individuals who are at risk for experiencing hemorrhagic episodes. In particularly preferred embodiments, the individuals are hemophiliacs with mild, moderate, or severe hemophilia.

As used herein, the term "subject" refers to any animal (*i.e.*, vertebrates and invertebrates), while the term "vertebrate subject" refers to any member of the subphylum Chordata. It is intended that the term encompass any member of this subphylum, including, but not limited to humans and other primates, rodents (*e.g.*, mice, rats, and guinea pigs), lagamorphs (*e.g.*, rabbits), bovines (*e.g.*, cattle), ovines (*e.g.*, sheep), caprines (*e.g.*, goats), porcines (*e.g.*, swine), equines (*e.g.*, horses), canines (*e.g.*, dogs), felines (*e.g.*, cats), domestic fowl (*e.g.*, chickens, turkeys, ducks, geese, other gallinaceous birds, etc.), as well as feral or wild animals, including, but not limited to, such animals as ungulates (*e.g.*, deer), bear, fish, lagamorphs, rodents, birds, etc. It is not intended that the term be limited to a particular age or sex. Thus,

adult and newborn subjects, as well as fetuses, whether male or female, are encompassed by the term.

As defined herein, a “therapeutically effective amount” or “therapeutic effective dose” is an amount or dose of AAV vector or virions capable of producing sufficient amounts of Factor VIII to decrease the time it takes for a subject’s blood to clot. Generally, severe hemophiliacs having less than 1% of normal levels of FVIII have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia A gene therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids encoding Factor VIII into a recipient host suspected of suffering from a blood clotting disorder. Using the nucleic acid as a template, the host produces Factor VIII, such that the subject’s blood is able to clot. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

I. AAV Vectors

Adeno-associated virus (AAV) is a non-pathogenic, replication-defective, helper-dependent parvovirus (or “dependovirus” or “adeno-satellite virus”). There are at least six recognized serotypes, designated as AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. Culture and serologic evidence indicates that human infection occurs with AAV-2 and AAV-3. Although 85% of the human population is seropositive for AAV-2, the virus has never been associated with disease in humans. Recombinant AAV (rAAV) virions are of interest as vectors for gene therapy because of their broad host range, excellent safety profile, and duration of transgene expression in infected hosts. One remarkable feature of recombinant AAV (rAAV) virions is the prolonged expression achieved after *in vivo* administration.

AAV vectors of the present invention may be constructed using known techniques to provide, as operatively linked components in the direction of transcription, (a) control sequences including a transcriptional initiation and termination regions, and (b) a nucleotide sequence encoding at least a portion of Factor VIII. The control sequences are selected to be functional in a targeted recipient cell. The resulting construct which contains the operatively linked components is bounded (5' and 3') with functional AAV ITR sequences.

The nucleotide sequences of AAV ITR regions are known (*See e.g.*, Kotin, Hum. Gene Ther., 5:793-801 [1994]; Berns, "Parvoviridae and Their Replication" in Fields and Knipe (eds), *Fundamental Virology*, 2nd Edition, for the AAV-2 sequence). AAV ITRs used in the vectors of the invention need not have a wild-type nucleotide sequence, and may be altered (*e.g.*, by the insertion, deletion or substitution of nucleotides). Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an AAV vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended.

A. Control Sequences

In some embodiments of the present invention, heterologous control sequences are employed with the vectors. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), a herpes simplex virus (HSV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE), a rous sarcoma virus (RSV) promoter, synthetic promoters, hybrid promoters, and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, also find use herein. Such promoter sequences are commercially available (*e.g.*, from Stratagene).

It is contemplated that in some embodiments, tissue-specific expression may be desirable (*e.g.*, expression of biologically active Factor VIII by hepatocytes). It is not intended that the present invention be limited to expression of biologically active Factor VIII by any particular cells or cell types. However, as hepatocytes (*i.e.*, liver cells) are the cells that normally synthesized Factor VIII (*See*, Kaufman, Ann. Rev. Med., 43:325 [1992]), it is contemplated that in some particularly preferred embodiments, the compositions of the present invention be administered to the liver.

In preferred embodiments, expression is achieved by coupling the coding sequence for Factor VIII with heterologous control sequences derived from genes that are specifically transcribed by a selected tissue type. A number of tissue-specific promoters have been described above which enable directed expression in selected tissue types. However, control sequences used in the present AAV vectors can also comprise control sequences normally associated with the selected nucleic acid sequences.

B. Construction of AAV Factor VIII Vectors

AAV vectors that contain a control sequence and a nucleotide sequence of interest (*i.e.*, at least a portion of the sequence encoding Factor VIII), bounded by AAV ITRs (*i.e.*, AAV vectors), can be constructed by directly inserting selected sequences into an AAV genome with the major AAV open reading frames ("ORFs") excised. Other portions of the AAV genome can also be deleted, so long as a sufficient portion of the ITRs remain to allow for replication and packaging functions. These constructs can be designed using techniques well known in the art (*See e.g.*, U.S. Patent Nos. 5,173,414 and 5,139,941, all of which are herein incorporated by reference); International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski *et al.*, Mol. Cell. Biol., 8:3988-3996 [1988]; Vincent *et al.*, Vaccines 90 [Cold Spring Harbor Laboratory Press, 1990]; Carter, Curr. Opin. Biotechnol., 3:533-539 [1992]; Muzyczka, Curr. Top. Microbiol. Immunol., 158:97-129 [1992]; Kotin, Hum. Gene

Ther., 5:793-801 [1994]; Shelling and Smith, Gene Ther., 1:165-169 [1994]; and Zhou *et al.*, J. Exp. Med., 179:1867-1875 [1994]).

Alternatively, AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct that is present in another vector using standard ligation techniques, such as
5 those described in Sambrook *et al.*, *supra*. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for
10 "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration). AAV vectors which contain ITRs have been described in (*e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference). In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC")
15 under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

Additionally, chimeric genes can be produced synthetically to include AAV ITR sequences arranged 5' and 3' of a selected nucleic acid sequence. The complete chimeric sequence is assembled from overlapping oligonucleotides prepared by standard methods (*See e.g.*, Edge, Nature 292:756 [1981]; Nambair *et al.*, Science
20 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Moreover, it is not intended that the present invention be limited to any specific Factor VIII sequence. Many natural and recombinant forms of Factor VIII have been isolated and assayed both *in vitro* and *in vivo*, using a variety of different regulatory elements and control sequences. Therefore, any known, or later discovered, DNA
25 sequence coding for biologically active Factor VIII can be expressed, alone or in combination with at least one additional vector, using the AAV vectors and methods taught in the present invention. Examples of naturally occurring and recombinant forms of Factor VIII can be found in the patent and scientific literature including, U.S. 5,563,045, U.S. 5,451,521, U.S. 5,422,260, U.S. 5,004,803, U.S. 4,757,006, U.S.

5,661,008, U.S. 5,789,203, U.S. 5,681,746, U.S. 5,595,886, U.S. 5,045,455, U.S.
5,668,108, U.S. 5,633,150, U.S. 5,693,499, U.S. 5,587,310, U.S. 5,171,844, U.S.
5,149,637, U.S. 5,112,950, U.S. 4,886,876, WO 94/11503, WO 87/07144, WO
92/16557, WO 91/09122, WO 97/03195, WO 96/21035, WO 91/07490, EP 0 672
5 138, EP 0 270 618, EP 0 182 448, EP 0 162 067, EP 0 786 474, EP 0 533 862,
EP 0 506 757, EP 0 874 057, EP 0 795 021, EP 0 670 332, EP 0 500 734, EP 0
232 112, EP 0 160 457, Sanberg *et al.*, XXth Int. Congress of the World Fed. Of
Hemophilia (1992), and Lind *et al.*, Eur. J. Biochem., 232:19 (1995).

Nucleic acid sequences coding for the above-described Factor VIII can be
10 obtained using recombinant methods, such as by screening cDNA and genomic
libraries from cells expressing Factor VIII or by deriving the sequence from a vector
known to include the same. Furthermore, the desired sequence can be isolated directly
from cells and tissues containing the same, using standard techniques, such as phenol
extraction and PCR of cDNA or genomic DNA (*See e.g.*, Sambrook *et al.*, *supra*, for a
15 description of techniques used to obtain and isolate DNA). Nucleotide sequences
encoding an antigen of interest (*i.e.*, Factor VIII sequence) can also be produced
synthetically, rather than cloned. The complete sequence can be assembled from
overlapping oligonucleotides prepared by standard methods and assembled into a
complete coding sequence (*See e.g.*, Edge, Nature 292:756 [1981]; Nambair *et al.*, Sci-
20 ence 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Although it is not intended that the present invention be limited to any
particular methods for assessing the production of biologically active Factor VIII, such
methods as immunoassays (*e.g.*, ELISA) and biological activity assays are
contemplated (*e.g.*, coagulation activity assays).

25 Furthermore, while in particularly preferred embodiments, human Factor VIII is
encompassed by the present invention, it is not intended that the present invention be
limited to human Factor VIII. Indeed, it is intended that the present invention
encompass Factor VIII from animals other than humans, including but not limited to
companion animals (*e.g.*, canines, felines, and equines), livestock (*e.g.*, bovines,

caprines, and ovines), laboratory animals (*e.g.*, rodents such as murines, as well as lagamorphs), and "exotic" animals (*e.g.*, marine mammals, large cats, etc.).

II. Virion Production

5 Producing AAV Factor VIII vectors and rAAV Factor VIII virions of the present invention generally involve the steps of: (1) introducing an AAV vector containing the Factor VIII gene into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions
10 missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions.

15 The above-described vectors and constructs can be introduced into a cell using standard methodology known to those of skill in the art (*e.g.*, transfection). A number of transfection techniques are generally known in the art (*See e.g.*, Graham *et al.*, Virol., 52:456 [1973], Sambrook *et al. supra*, Davis *et al., supra*, and Chu *et al.*, Gene 13:197 [1981]). Particularly suitable transfection methods include calcium phosphate
20 co-precipitation (Graham *et al.*, Virol., 52:456-467 [1973]), direct micro-injection into cultured cells (Capecchi, Cell 22:479-488 [1980]), electroporation (Shigekawa *et al.*, BioTechn., 6:742-751 [1988]), liposome-mediated gene transfer (Mannino *et al.*, BioTechn., 6:682-690 [1988]), lipid-mediated transduction (Felgner *et al.*, Proc. Natl. Acad. Sci. USA 84:7413-7417 [1987]), and nucleic acid delivery using high-velocity
25 microprojectiles (Klein *et al.*, Nature 327:70-73 [1987]).

 For the purposes of the invention, suitable host cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule. The term includes the progeny of the original cell which has been transfected. Thus, as

indicated above, a "host cell" as used herein generally refers to a cell which has been transfected with an exogenous DNA sequence. Cells from the stable human cell line, 293 (ATCC Accession No. CRL1573) are preferred in the practice of the present invention. Particularly, the human cell line 293 is a human embryonic kidney cell line that has been transformed with adenovirus type-5 DNA fragments (Graham *et al.*, J. Gen. Virol., 36:59 [1977]), and expresses the adenoviral E1a and E1b genes (Aiello *et al.*, Virol., 94:460 [1979]). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

Host cells containing the above-described AAV vectors must be rendered capable of providing AAV helper functions in order to replicate and encapsidate the nucleotide sequences flanked by the AAV ITRs to produce rAAV virions. AAV helper functions are generally AAV-derived coding sequences which can be expressed to provide AAV gene products that, in turn, function in *trans* for productive AAV replication. AAV helper functions are used herein to complement necessary AAV functions that are missing from the AAV vectors. Thus, AAV helper functions include one, or both of the major AAV ORFs, namely the *rep* and *cap* coding regions, or functional homologues thereof.

AAV helper functions are introduced into the host cell by transfecting the host cell with an AAV helper construct either prior to, or concurrently with, the transfection of the AAV vector. AAV helper constructs are thus used to provide at least transient expression of AAV *rep* and/or *cap* genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

In preferred embodiments, these constructs are in the form of a vector, including, but not limited to, plasmids, phages, transposons, cosmids, viruses, or virions. A number of AAV helper constructs have been described, such as the commonly used plasmids pAAV/Ad and pIM29+45 which encode both Rep and Cap expression products (See *e.g.*, Samulski *et al.*, J. Virol., 63:3822-3828 [1989]; and McCarty *et al.*, J. Virol., 65:2936-2945 [1991]). A number of other vectors have been

described which encode Rep and/or Cap expression products (*See e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference).

Both AAV vectors and AAV helper constructs can be constructed to contain one or more optional selectable markers. Suitable markers include genes which confer antibiotic resistance or sensitivity to, impart color to, or change the antigenic characteristics of those cells which have been transfected with a nucleic acid construct containing the selectable marker when the cells are grown in an appropriate selective medium. Several selectable marker genes that are useful in the practice of the invention include the gene encoding aminoglycoside phosphotransferase (APH) that allows selection in mammalian cells by conferring resistance to G418 (Sigma). Other suitable markers are known to those of skill in the art.

The host cell (or packaging cell) must also be rendered capable of providing non-AAV derived functions, or "accessory functions," in order to produce rAAV virions. Accessory functions are non-AAV derived viral and/or cellular functions upon which AAV is dependent for its replication. Thus, accessory functions include at least those non-AAV proteins and RNAs that are required in AAV replication, including those involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of *rep* and *cap* expression products and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses.

Particularly, accessory functions can be introduced into and then expressed in host cells using methods known to those of skill in the art. Commonly, accessory functions are provided by infection of the host cells with an unrelated helper virus. A number of suitable helper viruses are known, including adenoviruses; herpesviruses such as herpes simplex virus types 1 and 2; and vaccinia viruses. Nonviral accessory functions will also find use herein, such as those provided by cell synchronization using any of various known agents (*See e.g.*, Buller *et al.*, J. Virol., 40:241-247 [1981]; McPherson *et al.*, Virol., 147:217-222 [1985]; and Schlehofer *et al.*, Virol., 152:110-117 [1986]).

Alternatively, accessory functions can be provided using an accessory function vector. Accessory function vectors include nucleotide sequences that provide one or more accessory functions. An accessory function vector is capable of being introduced into a suitable host cell in order to support efficient AAV virion production in the host cell. Accessory function vectors can be in the form of a plasmid, phage, virus, transposon or cosmid. Accessory vectors can also be in the form of one or more linearized DNA or RNA fragments which, when associated with the appropriate control sequences and enzymes, can be transcribed or expressed in a host cell to provide accessory functions.

Nucleic acid sequences providing the accessory functions can be obtained from natural sources, such as from the genome of adenovirus, or constructed using recombinant or synthetic methods known in the art. In this regard, adenovirus-derived accessory functions have been widely studied, and a number of adenovirus genes involved in accessory functions have been identified and partially characterized (*See e.g.*, Carter, "Adeno-Associated Virus Helper Functions," in *CRC Handbook of Parvoviruses*, Vol. I (P. Tijssen, ed.) [1990], and Muzyczka, *Curr. Top. Microbiol. Immun.*, 158:97-129 [1992]). Specifically, early adenoviral gene regions E1a, E2a, E4, VAI RNA and, possibly, E1b are thought to participate in the accessory process (Janik *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1925-1929 [1981]). Herpesvirus-derived accessory functions have been described (*See e.g.*, Young *et al.*, *Prog. Med. Virol.*, 25:113 [1979]). Vaccinia virus-derived accessory functions have also been described (*See e.g.*, Carter, *supra.*, and Schlehofer *et al.*, *Virol.*, 152:110-117 [1986]).

As a consequence of the infection of the host cell with a helper virus, or transfection of the host cell with an accessory function vector, accessory functions are expressed which transactivate the AAV helper construct to produce AAV Rep and/or Cap proteins. The Rep expression products direct excision of the recombinant DNA (including the DNA of interest encoding at least a portion of Factor VIII) from the AAV vector. The Rep proteins also serve to duplicate the AAV genome. The expressed Cap proteins assemble into capsids, and the recombinant AAV genome is

packaged into the capsids. Thus, productive AAV replication ensues, and the DNA is packaged into rAAV virions.

Following recombinant AAV replication, rAAV virions can be purified from the host cell using a variety of conventional purification methods, such as CsCl
5 gradients. Further, if helper virus infection is employed to express the accessory functions, residual helper virus can be inactivated, using known methods. For example, adenovirus can be inactivated by heating to temperatures of approximately 60°C for approximately 20 minutes or more, as appropriate. This treatment selectively inactivates the helper adenovirus which is heat labile, while preserving the rAAV
10 which is heat stable.

III. Pharmaceutical Compositions

The resulting rAAV virions are then ready for use in pharmaceutical compositions which can be delivered to a subject, so as to allow production of
15 biologically active Factor VIII. Pharmaceutical compositions comprise sufficient genetic material that allows the recipient to produce a therapeutically effective amount of Factor VIII so as to reduce, stop and/or prevent hemorrhage. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible
20 pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, clotting factors or factor precursors, drugs or hormones. In preferred embodiments, the pharmaceutical compositions also contain a pharmaceutically acceptable excipient. Such excipients include any pharmaceutical
25 agent that does not itself induce an immune response harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol, sugars and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides,
30 hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such

as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., N.J. [1991]).

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art (*e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes).

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment. For administration of Factor VIII-containing vectors, such labeling would include amount, frequency, and method of administration.

5 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. Determining a therapeutic effective dose is well within the capability of those skilled in the art using the techniques taught in the present invention, such as ELISA and ChromZ FVIII coagulation activity assay, and other
10 techniques known in the art. Therapeutic doses will depend on, among other factors, the age and general condition of the subject, the severity of hemophilia, and the strength of the control sequences. Thus, a therapeutically effective amount in humans will fall in a relatively broad range that can be determined through clinical trials.

It is intended that the dosage treatment and regimen used with the present
15 invention will vary, depending upon the subject and the preparation to be used. Thus, the dosage treatment may be a single dose schedule or a multiple dose schedule. Moreover, the subject may be administered as many doses as appropriate to achieve or maintain the desired blood clotting time.

Direct delivery of the pharmaceutical compositions *in vivo* will generally be
20 accomplished via injection using a conventional syringe, although other delivery methods such as convention-enhanced delivery are envisioned (*See e.g.*, United States Patent No. 5,720,720, incorporated herein by reference). In this regard, the compositions can be delivered subcutaneously, epidermally, intradermally, intrathecally, intraorbitally, intramucosally (*e.g.*, nasally, rectally and vaginally),
25 intraperitoneally, intravenously, intraarterially, orally, or intramuscularly. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. In particularly preferred embodiments, the compositions are administered intravenously in the portal vasculature or hepatic artery

One skilled in the art will recognize that the methods and compositions
30 described above are also applicable to a range of other treatment regimens known in

the art. For example, the methods and compositions of the present invention are compatible with *ex vivo* therapy (e.g., where cells are removed from the body, incubated with the AAV vector and the treated cells are returned to the body).

IV. ADMINISTRATION

AAV vector can be administered to any tissue suitable for the expression of Factor VIII. In a preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal vasculature or hepatic artery where it is thought, without being bound by theory, that the vector transduces hepatocytes.

Current approaches to targeting genes to the liver have focused upon *ex vivo* gene therapy. *Ex vivo* liver-directed gene therapy involves the surgical removal of liver cells, transduction of the liver cells *in vitro* (e.g., infection of the explanted cells with recombinant retroviral vectors) followed by injection of the genetically modified liver cells into the liver or spleen of the patient. A serious drawback for *ex vivo* gene therapy of the liver is the fact that hepatocytes cannot be maintained and expanded in culture. Therefore, the success of *ex vivo* liver-directed gene therapy depends upon the ability to efficiently and stably engraft the genetically modified (*i.e.*, transduced) hepatocytes and their progeny. It has been reported that even under optimal conditions, autologous modified liver cells injected into the liver or spleen which engraft represent only a small percentage (less than 10%) of the total number of cells in the liver. Ectopic engraftment of transduced primary hepatocytes into the peritoneal cavity has been tried, in order to address the problem of engraftment in the liver.

Given the problems associated with *ex vivo* liver-directed gene therapy, *in vivo* approaches have been investigated for the transfer of genes into hepatocytes, including the use of recombinant retroviruses, recombinant adenoviruses, liposomes and molecular conjugates. While these *in vivo* approaches do not suffer from the drawbacks associated with *ex vivo* liver-directed gene therapy, they do not provide a means to specifically target hepatocytes. In addition, several of these approaches require performance of a partial hepatectomy, in order to achieve prolonged expression

of the transferred genes *in vivo*. Adenovirus and molecular conjugate based delivery methods also result in liver toxicity and inflammation which is an undesirable feature of Factor VIII gene therapy. The present invention provides compositions and methods for the long-term expression of biologically active Factor VIII. It is contemplated that the present invention will bypass the need for partial hepatectomy, while allowing expression of Factor VIII in concentrations that are therapeutic *in vivo*. The present invention further provides gene therapy compositions and methods that target hepatocytes for the production of Factor VIII by treated individuals.

Other tissues, however, may be suitable for the expression of Factor VIII even if they are not the tissue that normally synthesizes the protein. Muscle cells, for example, have been shown to express biologically active blood clotting Factor IX even though it is normally synthesized in the liver.

Finally, the AAV vectors may contain any nucleic acid sequences coding for biologically active Factor VIII. Additionally, the AAV vectors may contain a nucleic acid coding for fragments of Factor VIII which is itself not biologically active, yet when administered into the subject improves or restores the blood clotting time. For example, as discussed above, the Factor VIII protein comprises two polypeptide chains: a heavy chain and a light chain separated by a B-domain which is cleaved during processing. As demonstrated by the present invention, co-transducing recipient cells with the Factor VIII heavy and light chains leads to the expression of biologically active Factor VIII. Because, however, most hemophiliacs contain a mutation or deletion in only one of the chains (*e.g.*, heavy or light chain), it may be possible to administer only the chain defective in the patient and allow the patient to supply the other chain. In this case, the AAV vector would fall within the scope of the invention even though the single chain (*i.e.*, heavy or light) would not be biologically active until it was administered into a subject which can supply the second chain, thus forming biologically active Factor VIII.

V. FACTOR VIII ASSAYS

As described in the Experimental section below, there are many ways to assay Factor VIII expression and activity. Although the present invention is not limited to immunoassay methods, the present invention also provides methods for detecting Factor VIII expression comprising the steps of: a) providing a sample suspected of containing Factor VIII, and a control containing a known amount of known Factor VIII; and b) comparing the test sample with the known control, to determine the relative concentration of Factor VIII in the sample. Thus, the methods are capable of identifying samples (*e.g.*, patient samples) with sufficient or insufficient quantities of Factor VIII. In addition, the methods may be conducted using any suitable means to determine the relative concentration of Factor VIII in the test and control samples, including but not limited to means selected from the group consisting of Western blot analysis, Northern blot analysis, Southern blot analysis, denaturing polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), reverse transcriptase-coupled polymerase chain reaction (RT-PCR), enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescent immunoassay (IFA). Thus, the methods may be conducted to determine the presence of normal Factor VIII sequences in the genome of the animal source of the test sample, or the expression of Factor VIII (mRNA or protein), as well as detect the presence of abnormal or mutated Factor VIII gene sequences in the test samples.

In one preferred embodiment, the presence of Factor VIII is detected by immunochemical analysis. For example, the immunochemical analysis can comprise detecting binding of an antibody specific for an epitope of Factor VIII. In one another preferred embodiment of the method, the antibody comprises polyclonal antibodies, while in another preferred embodiment, the antibody comprises monoclonal antibodies.

It is further contemplated that antibodies directed against at least a portion of Factor VIII will be used in methods known in the art relating to the localization and structure of Factor VIII (*e.g.*, for Western blotting), measuring levels thereof in appropriate biological samples, etc. The antibodies can be used to detect Factor VIII in a biological sample from an individual (*e.g.*, an individual treated using the methods

and/or compositions of the present invention). The biological sample can be a biological fluid, including, but not limited to, blood, serum, plasma, interstitial fluid, urine, cerebrospinal fluid, synovial fluid, and the like. In particular, the antigen can be detected from cellular sources, including, but not limited to, hepatocytes. For
5 example, cells can be obtained from an individual and lysed (*e.g.*, by freeze-thaw cycling, or treatment with a mild cytolytic detergent including, but not limited to, TRITON X-100, digitonin, NONIDET P (NP)-40, saponin, and the like, or combinations thereof; *See, e.g.*, International Patent Publication WO 92/08981).

The biological samples can then be tested directly for the presence of the
10 Factor VIII using an appropriate strategy (*e.g.*, ELISA or RIA) and format (*e.g.*, microwells, dipstick [*e.g.*, as described in International Patent Publication WO 93/03367], etc.). Alternatively, proteins in the sample can be size separated (*e.g.*, by polyacrylamide gel electrophoresis (PAGE), with or without sodium dodecyl sulfate (SDS), and the presence of Factor VIII detected by immunoblotting [*e.g.*, Western
15 blotting]). Immunoblotting techniques are generally more effective with antibodies generated against a peptide corresponding to an epitope of a protein, and hence, are particularly suited to the present invention. In another preferred embodiment, the level of Factor VIII is assayed using the whole-blood clotting time and activated partial thromboplastin time (aPTT) of the subject's blood using techniques well known in the
20 art (Herzog *et al.*, Nature Medicine 5:56 [1999]).

The foregoing explanations of particular assay systems are presented herein for purposes of illustration only, in fulfillment of the duty to present an enabling disclosure of the invention. It is to be understood that the present invention contemplates a variety of immunochemical assay protocols within its spirit and scope.
25 Indeed, other methods such as biological assays to determine the presence and activity of Factor VIII are also encompassed by the present invention.

Thus, in addition to the immunoassay systems described above, other assay systems, such as those designed to measure and/or detect Fraction VIII and/or clotting

ability of a subject's blood are also encompassed by the present invention (*e.g.*, the ChromZ FVIII coagulation activity [FVIII-c] assay [Helena Labs]).

EXPERIMENTAL

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

In the experimental disclosure which follows, the following abbreviations apply: N (Normal); M (Molar); mM (millimolar); μ M (micromolar); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); mU (milliunits); ^{51}Cr (Chromium 51); μ Ci (microcurie); EC (degrees Centigrade); hFVIII (human factor VIII); FVIII (factor VIII); pH (hydrogen ion concentration); JRH grade; NaCl (sodium chloride); HCl (hydrochloric acid); OD (optical density); bp (base pair(s)); ATP (adenosine 5'-triphosphate); PCR (polymerase chain reaction); DNA (deoxyribonucleic acid); cDNA (complementary DNA); AAV (adeno-associated virus); rAAV (recombinant adeno-associated virus); ITR (inverted terminal repeat); FCS or FBS (fetal calf serum; fetal bovine serum); CFA (complete Freund's adjuvant); BSA (bovine serum albumin); ATCC (American Type Culture Collection, Rockville, MD); Sigma (Sigma Aldrich, St. Louis, MO); Biodesign International (Biodesign International, Kennebunk, MI); Baxter Hyland (Baxter Healthcare Corp., Biotech Group--Hyland Division, Hayward, CA); Helena Labs (Helena Laboratories, Beaumont, TX); American Diagnostica (American Diagnostica, Greenwich, CT); Accurate Chemical (Accurate Chemical and Scientific Corp., Westbury, NY); Molecular Probes (Molecular Probes, Eugene, OR); Vysis (Vysis, Downer Grove, IL); Tel-Test (Tel-Test, Inc., Friendswood, TX); Molecular Dynamics (Molecular Dynamics, Sunnyvale, CA); NUNC (Naperville, IL); and Stratagene

(Stratagene Cloning Systems, La Jolla, CA); Affinity Biologicals (Affinity Biologicals, Inc., Hamilton, Ontario); and Biodesign (Biodesign International, Kennebunkport, ME).

5

EXAMPLE 1

Dual Vector Plasmid Construction

The heavy and light chains of human Factor VIII (hFVIII) were assembled according to those reported by Yonemura *et al* (Yonemura *et al.*, Prot. Engineer., 6:669-674 [1993]) and cloned as expression cassettes into AAV vectors. Both vectors
10 contain the promoter and the first non-coding intron (from -573 to +985) from the human elongation factor 1 α (EF1 α) gene (Uetsuki *et al*, J. Biol. Chem., 264:5791-5798 [1989]; and Kim *et al.*, Gene 9:217-223 [1990]). Each vector also contains the first 57 base pairs of the FVIII heavy chain encoding the 19 amino acid signal sequence. The heavy chain construct encodes the A1 and A2 domains and 5 amino
15 acids from the N terminus of the B domain. The light chain vector encodes 85 amino acids of the carboxy terminal B domain, in addition to the A3, C1, and C2 domains. Both vectors utilize the human growth hormone (hGH) polyadenylation signal. The expression cassettes were inserted between AAV ITRs. The initial cloning step involved deleting 854 bp of EF1 α sequences between the *Spe*I and *Xcm*I sites of
20 pVm4.1e-hFIX (Nakai *et al.*, Blood 91:1-9 [1998]), and religating to create pVm4.1e δ D-hFIX.

This construct was then digested with *Eco*RI, which released the hFIX cDNA, and was ligated to an oligonucleotide containing *Mfe*I ends (*Eco*RI-compatible) and an internal *Cla*I restriction site, creating pVm4.1e δ D-linker. The heavy and light chain
25 fragments, including the hGH polyadenylation sequences were isolated from pVm4.1cFVIII-HC and pVm4.1cFVIII-LC, respectively as *Cla*I-*Bst*EII fragments. These fragments were cloned between the corresponding sites in the pVm4.1e δ D-linker, creating plasmids pVm4.1e δ D-FVIII-HC (also, rAAV-hFVIII-HC) and pVm4.1e δ D-FVIII-LC (also, rAAV-hFVIII-LC).

Figure 7 provides a map of the constructs. In this figure, the upper line in each panel represents the gene structure of the vectors, and the lower line represents the structure of the hFVIII protein domains encoded by the vectors (ITR, AAV inverted terminal repeat; EF1 α Pro/Intron 1, human polypeptide elongation factor 1 α gene promoter and first intron; hFVIII-HC human FVIII cDNA; hFVIII-LC, human FVIII cDNA; hGH PA, human growth hormone polyadenylation signal; SS, human FVIII signal sequence; A1, A2, "B", A3, C1, C2, complete and incomplete ("") protein domains of the hFVIII protein).

EXAMPLE 2

Single Vector Plasmid Construction

The plasmid pAAV-F8-1 construct containing both the light and heavy chains of factor VIII was constructed as follows. A PCR fragment, Z8, containing cloning sites, 5' -splicing donor site of a synthetic intron based on EF1 α and immunoglobulin G (IgG) intron sequences, Kozak sequence and the first 16 nucleotides of the human blood coagulation factor VIII (FVIII) coding sequence was generated using oligonucleotides Z8S and Z8A. The sequences of the nucleic acids is shown below:

Oligonucleotide Z8S:

5' cccaagcttgcggccgccgggtgccgcccctaggcaggtaagtgccgtgtgtggtcc 3'
(SEQ ID NO:1)

Oligonucleotide Z8A:

5' ccgctcgagcagagctctatttgcattggtggaatcgatgccgcgggaaccacacacggc 3'
(SEQ ID NO:2)

PCR fragment Z8:

5' cccaagcttgcggccgccgggtgccgcccctaggcaggtaagtgccgtgtgtggtcccgccgctcgcgagcgg 3' (SEQ ID NO:3)

Nucleic acid Z8 was inserted into pZERO-2 (Invitrogen) between *HindIII* and *XhoI* sites to create pZ8. A PCR fragment, INT3, containing the branching point, the

polypyrimidine tract, and the 3' splicing acceptor site of the synthetic intron was generated using oligonucleotides INT3S and INT3A whose sequence is shown below.

Oligonucleotide INT3S:

5' ttcccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactga 3'
(SEQ ID NO:4)

Oligonucleotide INT3A:

5' gaatcgatacctgtggagaaaaagaaaaagtggtatgcagtcagtaattcaaggc 3'
(SEQ ID NO:5)

PCR Fragment INT3:

5' ttcccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactgacact
gacatccactttttcttttccacaggtatcgattc 3' (SEQ ID NO:6)

15

INT3 was inserted between the *Sac*II and *Cla*I sites of pZ8 to create pZ8.I. Therefore, Z8.I contains the entire synthetic intron between *Avr*II and *Cla*I sites. A hFVIII cDNA fragment having *Sac*I and *Xho*I restriction sites was inserted between the *Sac*I and *Xho*I sites of pZ8.I to create pZ8.I.dB. Therefore, pZ8.I.dB contains a synthetic intron and the entire coding sequence of hFVIII.

20

A PCR fragment, EG3, containing three HNF-3 binding sites and -54 to +8 of mouse albumin gene was generated using oligonucleotides EG3S and EG3A with modifications to eliminate linker sequences. The sequences of EG3S and EG3A are as follows:

25

Oligonucleotide EG3S:

5' aggggaatgtttgttcttaataaccatccaggggaatgtttgttcttaataaccatccaggggaatgtttgttcttaatacca
tctacagtattgtgtaaa 3' (SEQ ID NO:7)

30

Oligonucleotide EG3A:

5' ggaaaggtgatctgtgtgcagaaagactcgctctaatacttctttaaccaataactg 3'
(SEQ ID NO:8)

5 PCR Fragment EG3:

5' aggggaatgtttgttcttaataaccatccaggggaatgtttgttcttaataaccatccaggggaatgtttgttctta
aataccatctacagtatttggttaaagaagtatattagagcgagtccttctgcacacagatcacctttcc 3'
(SEQ ID NO:9)

10 EG3 was then phosphorylated using T4 polynucleotide kinase and inserted into
the *Sma*I site of pZ8.I.dB to create pZ8.I.dB.egg. A DNA fragment, SPA, containing
an efficient synthetic polyA signal based on rabbit β -globin sequences (Genes and
Develop., 3:1019) was generated by hybridizing two oligonucleotides SPA.S and
SPA.A.

15

Oligonucleotide SPA.S:

5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgc 3'
(SEQ ID NO:10)

20 Oligonucleotide SPA.A:

5' tcgagcggccgcacacaaaaaccaacacacagatctctagagctctgatctttttatc 3'
(SEQ ID NO:11)

PCR Fragment SPA:

25 5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgctcga 3'
(SEQ ID NO:12)

SPA was inserted into the *Xho*I site of pZero-2 to create pZero-2.SPA. SPA
was excised from a pZero-2.SPA clone and inserted into the *Xho*I site of pZ8.I.dB.egg

to create pZ8.I.dB.egg.A. pAAV-CMV-FIX9 was digested with *Cla*I, blunted with T4 polymerase and religated to create pAAV(*Cla*-)-CMV-FIX9.

The entire expression cassette containing HNF-3.albumin promoter-synthetic intron-hFVIII-synthetic poly A signal was excised from pZ8.I.dB.egg.A using *Not*I and
5 ligated to the plasmid backbone and AAV ITRs from pAAV (*Cla*-)-CMV-FIX9 to create pAAV-F8-1. The nucleotide sequence of the vector from ITR to ITR (*i.e.*, excluding plasmid backbone) is shown in SEQ ID NO 13.

EXAMPLE 3

Virion Production

AAV vectors were produced from these plasmids using the Ad free system as previously described in U.S. Patent No. 5,858,351; U.S. Patent No. 5,846,528; U.S. Patent No. 5,622,856; and Matsushita *et al.*, Gene Ther 5:938 (1998) all of which are hereby incorporated by reference. Briefly, 293 cells (ATCC, catalog number CRL-
15 1573) were seeded in 10 cm dishes at a density of 3×10^6 cells per dish in 10 ml medium and incubated at 37°C with CO₂ and humidity. After an overnight incubation, cells were approximately seventy to eighty percent confluent.

The cells were then transfected with DNA by the calcium phosphate method, as is well known in the art. Briefly, 7 µg of AAV vector containing the Factor VIII
20 coding region, 7 µg of pladenos which supplies the accessory functions, and 7 µg of 1909 AAV helper were added to a 3 ml sterile, polystyrene snap cap tube using sterile pipette tips. Then, 1.0 ml of 300 mM CaCl₂ (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na₂PO₄, 12 mM dextrose, pH 7.05, JRH grade)
25 was added with a 2 ml pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the cells, one drop at a time, evenly throughout the 10 cm dish. The cells were then incubated at 37°C with CO₂ and humidity for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were
30 provided with fresh medium and incubated for 72 hours.

After 72 hours, the cells were harvested, pelleted, and resuspended in 1 ml TBS/1% BSA. Freeze/thaw extracts were prepared by repeatedly (three times) freezing the cell suspension on dry ice and thawing at 37°C. Viral preps were stored at -80°C and titered by dot blot assay prior to the first round of infection.

5

EXAMPLE 4

In Vitro Cell Transduction

Cells from the stable human cell line, 293 (ATCC No. CRL1573) were seeded in six-well plates (*i.e.*, plates having six wells for cell growth) at a density of 5×10^5 cells/well. When the monolayers reached 80-90% confluence, they were infected with rAAV virions AAV-e δ D-FVIII-HC, AAV-e δ D-FVIII-LC, an equal ratio of AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC, or AAV-e δ D-FIX at MOIs of 3×10^3 and 3×10^4 . Eighteen hours post infection, the media were replaced with DMEM/10% heat inactivated FBS. The media were collected later for analysis by ELISA (as described below) for FVIII light chain antigen levels, and by the ChromZ FVIII as coagulation activity (FVIII-c) assay (Helena Labs) for biological activity, using the manufacturer's instructions and as described in Example 6.

EXAMPLE 5

Single Chain Factor VIII Infectivity Assay

In this Example, the infectivity of single chain Factor VIII was investigated. To determine the infectivity of rAAV-hF8-1, HepG2, 293, and H2.35 cells were infected with rAAV-hF8-1 and a control vector rAAV-hF8L at an MOI of 1×10^4 viral particles per cell. Recombinant AAV DNA in infected cells was isolated by Hirt extraction and run on an alkaline agarose gel. Southern blot analysis using an human F8 probe showed that similar amounts of rAAV-hF8-1 and rAAV-hF8L were isolated from uncoated virus in the infected cells. An infectious center assay (ICA) known in the art (*See e.g.*, Snyder, *Current Protocols in Genetics*, Chapter 12, John Wiley & Sons [1997]) was used to further characterize the infectivity of rAAV-hF8-1. In this assay, the infectious particle to total particle ratio of rAAV-F8-1 and that of a control

rAAV vector with the genome size of 4645 nucleotides was determined. The results indicated that rAAV-hF8-1 had an infectious particle to total particle ratio that was comparable to the control vector at approximately 1:1000. Taken together, these results indicate rAAV-hF8-1 has similar infectivity as rAAV vectors with the genome size of wild-type AAV.

EXAMPLE 6

Factor VIII Protein Expression Assay

An ELISA specific for the light chain of FVIII was used to determine FVIII light chain antigen levels in the 293 cells, as well as the injected animals (described below). NUNC Maxisorb 96 well plates were coated with 50 μ l of a 1:500 dilution of the light chain specific antibody, N77110 (Biodesign International) in a coating buffer overnight at 4°C. The plate was washed three times with wash buffer (PBS, 0.05% Tween 20) and blocked with 200 μ l blocking buffer (PBS, 10% horse serum, 0.05% Tween 20) at room temperature for 1 hour. The plate was washed three times and standards and samples were applied. Bioclone recombinant human FVIII (Baxter Hyland) was used as the standard, and was diluted in blocking buffer to concentrations ranging from 320 ng/ml to 10 ng/ml.

For analysis of transduced culture supernatants, the standards contained 50% media, and for analysis of mouse plasma, the standards were diluted into 10%, in normal pooled mouse plasma (Sigma). A standard assay reference plasma (SARP; Helena Labs) was also included in the assay. Following the loading of the standards and samples (95 μ l/well), the plate was incubated at room temperature for 2 hours, and washed five times with wash buffer (200 μ l/well). A 1:200 dilution of a horseradish peroxidase-conjugated light chain specific antibody, ESH8-HRP, (American Diagnostica) was added (100 μ l/well), and the plate was incubated for 1 hour at room temperature. The plates were then washed four times with wash buffer, and the antigen was detected using an ABTS peroxidase substrate kit (BioRad) according to the manufacturer's instructions. The results are shown in Table 1 of Example 7, below.

EXAMPLE 7

Factor VIII Biological Activity Assay

The ChromZ FVIII coagulation activity (FVIII-c) assay (Helena Labs, Beaumont, TX) was used to detect biologically active FVIII in the 293 cells infected as described in Example 4. Bioclone recombinant human FVIII (Baxter Hyland) was used as a standard to analyze transfected culture supernatants. The standards were diluted in plasma dilution buffer (supplied in kit) in the range of 10 ng/ml to 0.313 ng/ml, and were made 2.5% in media. Because this assay can detect both human and murine FVIII activity, it was modified to deplete biologically active human Factor VIII in the mouse plasma. Mouse plasma was pre-incubated with an antibody specific for human FVIII prior to performing the assay. The difference in FVIII activity between the untreated plasma sample and the antibody treated sample represent the amount of biologically active human FVIII in the plasma. The standard used in the assay was normal pooled human plasma (FACT; obtained from George King Biomedical). Serial dilutions of FACT were made in FVIII deficient plasma from undiluted (200 ng/ml) to 6.25 ng/ml. The standards (10 μ l) were incubated at 37°C for 15 min., with or without the addition of 2 μ l antibody N77110. Similarly, mouse plasma samples were diluted in FVIII deficient plasma and 10 μ l of these diluted samples were incubated with or without 2 μ l of N77110 at 37°C for 15 min., and immediately placed on ice. Thus, all incubations with antibody were done in a background of 100% plasma. The antibody adsorbed and non-adsorbed FACT standards, as well as the mouse plasma samples were diluted 1:20 in plasma detection buffer provided in the ChromZ kit. Thus, the final concentration of the FACT standards used in the assay ranged from 10 ng/ml to 0.313 ng/ml.

Twenty five microliters of these dilutions were added to a chilled 96 well plate. With the plate on ice, 25 μ l of FIXa reagent and 50 μ l of FX were added, and the plate was incubated at 37°C for 15 min. Substrate (50 μ l) was added and the plate was incubated for an additional 3 min at 37°C. The reaction was stopped with the addition of 25 μ l 50% acetic acid and the optical density at 405 nm was measured.

As shown below in Table 1, infection of 293 cells with AAV-e δ D-FVIII-HC resulted in no antigen production, as well as no biologically-active protein. Cells infected with AAV-e δ D-FVIII-LC produced FVIII light chain, but no biologically active protein. However, cells transduced with both vectors produced FVIII light chain and biologically active FVIII in a dose-dependent manner. Transduction of cells with the negative control vector, AAV-e δ D-FIX, resulted in no antigen nor any biologically active FVIII. It was assumed that equal amounts of heavy and light chains were produced in transduced cells. The activity units were converted to nanograms using the definition of one unit being equal to the amount of FVIII in 1 ml of normal pooled human plasma, or 200 ng.

**TABLE 1. *In Vitro* Production of Biologically Active
Human Factor VIII From Two rAAV Vectors**

	Vector	MOI	ELISA (ng/ml)	ChromZ	
				(mU/ml)	(ng/ml)
5	AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC	3×10^3	24	35	7.1
	AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC	3×10^4	121	440	87.9
	AAV-e8D-FVIII-HC	3×10^3	0	0	0
10	AAV-e8D-FVIII-HC	3×10^4	0	0	0
	AAV-e8D-FVIII-LC	3×10^3	20.5	0	0
	AAV-e8D-FVIII-LC	3×10^4	96.9	0	0
	AAV-hFIX	3×10^3	0	0	0
	AAV-hFIX	3×10^4	0	0	0
15	No Vector		0	0	0

EXAMPLE 8

Immunofluorescent Staining of FVIII Heavy and Light Chains

20 In these experiments, 293 cells transduced as described above were analyzed using immunofluorescent staining. 293 cells were plated on rat tail collagen-coated two-well culture slides at a density of 4×10^5 cells per well. Forty-eight hours later, the cells were transduced at an MOI of 3×10^4 particles per cell of rAAV-hFVIII-HC and rAAV-hFVIII-LC. Forty-eight hours post-transduction, the cells were fixed *in situ*

25 with acetone, blocked with 2% BSA, and stained with a fluorescently labelled anti-hFVIII light chain antibody and a fluorescently labelled anti-hFVIII heavy chain antibody. The anti-hFVIII light chain antibody used was ESH-4 monoclonal antibody (American Diagnostica), fluorescently labelled with alexa-488 (Molecular Probes),

according to the manufacturer's instructions. The anti-hFVIII heavy chain antibody used was MAS530P monoclonal antibody (Accurate Chemical) fluorescently labelled with alexa-594 (Molecular Probes), according to the manufacturer's instructions. The cells were counter-stained with DAPI. The images were collected using a Zeiss
5 Axioskop fluorescence microscope equipped with separate filters for DAPI, FITC, and rhodamine signals and a CCD camera. Image analysis was performed using Quips imaging software (Vysis).

As indicated above, infection of cells with either rAAV-e8D-FVIII-HC or AAV-e8D-FVIII-LC, followed by staining with antibodies to both chains resulted in
10 production of the individual chains of human FVIII. Immunofluorescent staining of cells co-infected with both vectors demonstrated that although some cells express only the heavy or light chain of hFVIII, many co-expressed both chains of human FVIII.

EXAMPLE 9

15 *In Vitro* Expression of Factor VIII Using Single Construct

Table 2 shows that two single vector constructs containing the heavy and the light chain of Factor VIII driven by different promoters express biologically active Factor VIII. The constructs pAAV-hF8-1 (SEQ ID NO:13), and pVm4.1cF8ΔB (SEQ ID NO:14) were transfected into 293 cells. Following transfection, the cells were
20 allowed to express factor VIII for 48-72 hours. Factor VIII in the culture media was assayed by the ChromZ FVIII coagulation activity (FVIII-c) assay, as per the manufacturer's instructions.

TABLE 2***In Vitro* Production of Biologically Active FVIII**

Construct(s)	ELISA (ng/ml)	ChromZ (ng/ml)
Control	-	0
pAAV-hF8-1	-	4.9
pVm4.1cF8ΔB	-	46

EXAMPLE 10**Factor VIII Expression Using Tissue Specific Promoters**

In these experiments, different promoters and enhancer elements were used to drive expression of a Factor VII coding sequence. Expression of Factor VIII was compared in 293 cells and HepG2 cells using different promoters. The pAAVeF8ΔB contains an EF-1 α promoter with a hGH intron, Factor VIII with a B-domain deletion (F8ΔB) and a polyA. As described previously, pAAV-hF8-1 uses the HNF-3 albumin promoter with a minimal intron followed by F8ΔB and a minimal polyA. The construct pAAV-c8 uses the CMV enhancer-promoter and the F8ΔB. pAAV8b1 contains the HNF-3 albumin promoter followed by the CMV/B-globin intron with the F8ΔB and a minimal poly A site. Table 3 describes Factor VIII expression using the albumin promoter relative to the control plasmid pV4.1cF8ΔB in HepG2 and 293 cells. These data show increased expression of Factor VIII in HepG2 liver cells with the albumin promoter as compared to Factor VIII expression in 293 cells.

TABLE 3

Relative Tissue Specificity of Promoters

Plasmid Construct	HepG2 Cells	293 Cells
pAAV-hF8-1	6.2	0.6
pAAV8bl	6.7	1.0
pAAVc8	30.0	41.0
pV4.1eF8ΔB	100	100

5

Next, several promoters derived from the transthyretin (TTR) gene promoter were transfected into HepG2 cells. TTR is an abundant serum protein and the gene enhancer-promoter contains well known liver-specific transcription factor binding sites (Samadani *et al.*, Gene Expression 6:23 [1996]; Yan *et al.*, EMBO 9:869 [1990]; Costa and Grayson, Nuc. Acids Res., 19:4139 [1991]; Costa *et al.*, Mol. Cell. Bio., 6:4697 [1986]). The constructs were made by replacing the HNF-3 albumin promoter in pAAV-hF8-1 with various lengths of the TTR promoter-enhancer. The TTR enhancer-promoter was modified by replacing the weak affinity binding sites with the strong affinity binding sites to create pAAV-hF8-2. The pAAV-hF8-TTR-E-L-P202 construct contains the full TTR promoter with a linker between the enhancer and the promoter. The remaining constructs are 5' deletions: pAAV-hF8-TTR-E-P202 has the promoter and enhancer with no linker; pAAV-hF8-TTR-E-P197 has a 5 base pair deletion from the promoter; pAAV-hF8-TTR-E-P151 has a 50 base pair deletion; pAAV-hF8-TTR-P202 lacks the TTR enhancer and pAAV-hF8-TTR(X) has a 65 base pair deletion in the enhancer. The control plasmid, pAAV-hF8-1, expressed approximately 4.6 mU/ml. Table 4 shows the fold-increase in Factor VIII activity using the TTR promoter series relative to the control plasmid.

15

20

25

TABLE 4

Factor VIII Expression Using TTR-Derived Promoters

Plasmid Construct	Relative Factor VIII Activity
pAAV-hF8-STTR	3.16
pAAV-hF8-TTR-E-L-P202	8.86
pAAV-hF8-TTR-E-P202	6.1
pAAV-hF8-TTR-EP197	7.3
pAAV-hF8-TTR-E-P151	13.3
pAAV-hF8-TTR-P202	2.3

EXAMPLE 11

***In Vivo* Expression of Factor VIII**

In order to test the feasibility of the AAV vector approach of the present invention *in vivo*, three groups of five C57BL/6 mice were injected via the portal vein with either 3×10^{11} particles of AAV-e δ D-FVIII-HC, 3×10^{11} particles of AAV-e δ D-FVIII-LC, or 3×10^{11} particles of both AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC. In addition, a group of four animals was injected with 3×10^{11} particles of AAV-e δ D-FIX. It has been shown that this strain of mice does not elicit an immune response to human FVIII when the gene is delivered to the liver via an adenoviral vector (Connelly *et al.*, Blood 87:4671-4677 [1996]). As indicated by the results shown below, the data obtained during these experiments demonstrate the feasibility of producing biologically active FVIII using two AAV vectors to independently deliver the heavy and light chains of FVIII.

Blood samples were collected in sodium citrate via the retro-orbital plexus at biweekly intervals for the first 2 months and at monthly intervals thereafter for 6

months and at 11 months. Very high levels of FVIII light chain were expressed in animals injected with AAV-e8D-FVIII-LC alone or both vectors as shown in Figure 8.

In order to assess the amount of biologically active human FVIII produced in the animals, a modified ChromZ assay was used. Since this assay detects both human and murine FVIII, the amount of FVIII present in the plasma before and after adsorption to an antibody specific to human FVIII was determined. The amount of FVIII remaining in the plasma after adsorption represented the amount of active murine FVIII and the difference represented the amount of active human FVIII. Control experiments demonstrated that the antibody could remove 80-90% of the human FVIII from a mouse plasma sample when the sample was spiked with up to 32 ng of human FVIII. The modified ChromZ assay indicated that only those animals injected with both vectors produced biologically active FVIII, as indicated in Table 5. The results shown in Table 5 are those from plasma collected 8 weeks post-injection, although similar results were obtained at 10 weeks and 5 months post-injection. One of the five animals co-injected with both the heavy and light chain vector did not express VIII, presumably due to an inefficient injection, and was omitted from the analysis. Animals injected with both vectors produced over 2 µg/ml hFVIII light chain as measured by ELISA. The ChromZ assay indicated that a total of 600-900 ng/ml of active hFVIII was detected in the plasma. The contribution from murine Factor VIII was approximately 400-500 ng/ml, indicating that about 230-430 ng/ml of active human Factor VIII was present in the plasma. Although only a fraction of the total protein was found to be active, the animals produced physiological levels of the active protein (*i.e.*, 200 ng/ml). The animals were found to have maintained these physiological levels of active protein for more than 11 months, without waning. Similar analyses performed on animals injected with the light chain vector alone, the heavy chain vector alone, or the hFIX vector demonstrated no biologically active human FVIII in the plasma of these animals.

TABLE 5

Biological Activity of Human Factor VIII *In Vivo*

Construct(s) Used	ELISA (ng/ml)	Total FVIII (-Ab) (Units)	Murine FVIII (+Ab) (Units)	Human FVIII (ng/ml)
AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC*	2288	3.9	2.2	342
AAV-e δ D-FVIII-LC *	3329	1.4	1.6	0
AAV-e δ D-FVIII-HC	0	1.6	1.6	0
AAV-e δ D-FIX	0	1.4	2.0	0

*Average of three animals.

EXAMPLE 12**Gene Transfer and Vector Expression in Tissues**

In these experiments, evidence of gene transfer to liver was obtained by Southern Blot analysis following isolation of DNA from one animal of each experimental group sacrificed 8 weeks post-injection (*i.e.*, as described in Example 11). In addition, DNA was obtained from other tissues in order to determine the degree of vector expression in organs other than the liver.

Twenty micrograms of DNA was digested with *Bgl*II, separated using a 1% agarose gel, and hybridized with a ³²P-labelled 1126 bp *Alw*NI fragment encoding the A1 and A2 domains of hFVIII (heavy chain probe), or a ³²P-labelled 1456 bp *Nde*I-*Eco*RI fragment encoding the A3, C1 and C2 domains of hFVIII (light chain probe).

Copy number controls were generated by spiking *Bgl*II-digested naive mouse liver DNA with *Bgl*II-digested heavy or light chain plasmid DNA (pVm4.1e δ D-hFVIII-HC and pVm4.1e δ D-hFVIII-LC, respectively), at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome. The hybridized membranes were analyzed using a Storm 860 phosphoimager (Molecular Dynamics), and quantitation of vector copy number was evaluated using ImageQuaNT software (Molecular Dynamics). Autoradiography of the

hybridized membranes was also performed. Total RNA was isolated from liver tissue using the RNA Stat extraction kit (Tel-Test). As describe briefly below, Northern blot analysis was also performed on 10 µg RNA using methods known in the art, in conjunction with the ³²P-labelled probes specific to the heavy and light chains of hFVIII described above and autoradiography was performed on the hybridized membranes.

Following digestion of liver DNA with *Bgl*II and hybridization with an hFVIII light chain probe described below, using methods known in the art, a band at the predicted size of 3015 bp was detected in animals injected with rAAV-hFVIII-LC, or both the heavy and light chain vectors. This band was not observed in the DNA of animals injected with the heavy chain vector alone or the hFIX vector, as shown in Figure 9A (rAAV-hFVIII-LC, lane 1; rAAV-hFIX, lane 2; rAAV-hFVIII-HC, lane 3; both rAAV-hFVIII-LC and rAAV-hFVIII-HC, lane 4; copy number controls were generated by spiking *Bgl*II digested naive mouse liver DNA with the corresponding plasmids at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome, lanes 5-9). Phosphoimage analysis revealed that the light chain vector was present at approximately 2.4 and 1.5 copies per diploid genome in animals injected with the light chain vector alone or both vectors, respectively. When *Bgl*II-digested DNA was hybridized with an hFVIII heavy chain probe, the expected band of 2318 bp was observed in animals injected with the heavy chain vector alone or both vectors, but was not detected in animals injected with the light chain vector alone or the hFIX vector, as shown in Figure 9B. The copy number in animals injected with the heavy chain vector alone and both vectors was 1.1 and 1.7 vector copies per diploid genome, respectively.

The results of hybridization of DNA extracted from the spleen, kidney and heart tissue with either an hFVIII light chain probe or a heavy chain probe indicated that these tissues contained less than 1 copy of vector sequences per 10 diploid genomes, demonstrating that the vector distributes primarily to the liver following intra-portal injection, as shown in Figures 10A and 10B.

Human FVIII gene expression in the liver of the mice was also assessed by Northern blot analysis on RNA isolated from animals sacrificed 8 weeks post-injection (as described above). hFVIII light chain transcripts of the predicted size (2.7 kb) were observed in animals injected with the light chain vector alone or both vectors, as
5 shown in Figure 11A. Similarly, the expected hFVIII heavy chain transcripts (2.7 kb) were detected in animals that were injected with the heavy chain vector alone or both vectors, as shown in Figure 11B. Since the heavy and light chain DNA sequences were shown by Southern blot analysis to be present at approximately the same copy number (1.7 and 1.5 copies per diploid genome, respectively), in an animal injected
10 with both vectors, these results demonstrate that both the heavy and light chains of hFVIII are expressed in the liver in approximately equivalent amounts.

EXAMPLE 13

15 *In vivo* Expression of Factor VIII Using Single Vectors

Several groups comprising four C57BL/6 mice were injected via the portal vein with 3×10^{11} particles of AAV-hF8-STTR, 3×10^{11} particles of AAV-hF8-hAAT-137, or 3×10^{11} particles of AAV-hF8-HNF3-alb-original. As discussed above, AAV-hF8-STTR contains modified sequences from the transthyretin gene promoter. AAV-hF8-
20 hAAT-137 contains 137 base pairs of the human₁-antitrypsin promoter. *See* De Simone et al., EMBO J., 6:2759-2766 (1987). AAV-hF8-HNF3-alb-original, like AAV-F8-1, contains three HNF3 binding sites and 54 base pairs of the albumin promoter.

Expression of Factor VIII was measured at 4 weeks by human Factor VIII-
25 specific ELISA (Affinity Biologicals). Even at this early time-point, several of these mice expressed between 2 and 20 ng/ml of human Factor VIII. Table 6 shows expression levels at 4 weeks post-infection for selected animals.

TABLE 6. Factor VIII Expression at 4 Weeks

rAAV	Factor VIII Levels (ng/ml)
rAAV-hF8-STTR (mouse #1)	4.3
rAAV-hF8-STTR (mouse #2)	2.4
rAAV-hF8-hAAT-137 (mouse #1)	10.5
rAAV-hF8-hAAT-137 (mouse #2)	20.4
rAAV-hF8-HNF3-alb-original (mouse #1)	2.2

10 rAAV-hF8-hAAT-137 produced expression levels of as much as 10 to 20 ng/ml, which represent 20% to 40% of normal human levels of Factor VIII. These data therefore show the *in vivo* expression of therapeutically effective amounts of human Factor VIII using single factor AAV vectors.

FIGURE 1




FIGURE 2




FIGURE 3

AAV-F8-1

-S-F-S-Q-N-P-P-V-L-K-R-H-Q-R-




FIGURE 4

PVM4.1c-F8AB

-S-F-S-Q-N-P-P-V-L-K-R-H-Q-R-




FIGURE 5

CAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCCGGCGCTCGGGCGACCTTTGGTTCGCCCCGGCCTCAGT
GAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGCCCGCCAGGGAATGTTTGTCTT
AAATACCATCCAGGGAATGTTTGTCTTAAATACCATCCAGGGAATGTTTGTCTTAAATACCATCTACAGTTATTGGTT
AAAGAAGTATATTAGAGCGAGTCTTTCTGCACACAGATCACCTTTCCGGGTGCCGCCCTAGGCAGGTAAGTGCCGTGTG
TGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTGCGTGCCTTGAATTACTGACACTGACATCCACTTTTTCT
TTTTCTCCACAGGTATCGATTCCACCATGCAAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGGGATTCTGCTTT
AGTGCCACCAGAAGATACTACCTGGGTGCAGTGGAACTGTCTATGGGACTATATGCAAAGTGATCTCGGTGAGCTGCCTGT
GGACGCAAGATTTCTCCTAGAGTGCCAAAATCTTTTCCATTCAACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAG
AATTCACGGATCACCTTTTCAACATCGCTAAGCCAAGGCCACCTGGATGGGTCTGCTAGGTCTACCATCCAGGCTGAG
GTTTATGATACAGTGGTCATTACACTTAAGAACATGGCTTCCCATCCTGTCACTCTTCATGCTGTTGGTGTATCCTACTG
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAAAAGATGATAAAGTCTTCCCTGGTGGAA
GCCATACATATGTCTGGCAGTCTGAAAGAGAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATATCTT
TCTCATGTGGACCTGGTAAAGACTTGAATTCAGGCCTCATTTGAGCCCTACTAGTATGTAGAGAAGGGAGTCTGGCCAA
GAAAAGACACAGACCTTGCACAAATTTATACTACTTTTTGCTGTATTTGATGAAGGAAAAGTTGGCACTCAGAAAACAA
AGAACTCCTTGATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTGGCCTAAAATGCACACAGTCAATGGTTATGTAAAC
AGGTCTCTGCCAGGTCTGATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTGGAATGGGCACCACCTCTGAAGT
GCACTCAATATTCTCGAAGGTACACATTTCTGTGAGGAACCATCGCCAGGCGTCTTGGAATCTCGCCAATAACTT
TCCTTACTGCTCAAACTCTTGATGGACCTTGGACAGTTTCTACTGTTTTGTGCATATCTCTTCCCACCAACATGATGGC
ATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCCCACTACGAATGAAAAATAATGAAGAAGCGGAAGACTA
TGATGATGATCTTACTGATTCTGAAATGGATGTGGTCAAGTTTGTATGATGACAACCTCTCCTTCTTTATCCAAATTCGCT
CAGTTGCCAAGAAGCATCTAAAACCTTGGGTACATTACATTGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTC
CTCGCCCCGATGACAGAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAGGAAGTACAAAAAGT
CCGATTTATGGCATAACAGATGAAACCTTTAAGACTCGTGAAGCTATTTCAGCATGAATCAGGAATCTTGGACCTTTAC
TTTATGGGGAAGTTGGAGACACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATATAACATCTACCCCTCACGGA
ATCACTGATGTCCGTCTTTGTATTCAAGGAGATTACAAAAGGTGTAAAACATTTGAAGGATTTTCCAATTTCTGCCAGG
AGAAATATTCAAATATAAATGGACAGTGAAGTGTAGAAGATGGGCCAACTAAATCAGATCCTCGGTGCCGTGACCCGCTATT
ACTCTAGTTTCGTTAATATGGAGAGAGATCTAGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTA
GATCAAAGAGGAAACCAGATAATGTGACAGAAGAGGAATGTCTCTGTTTTCTGTATTTGATGAGAACCAGGCTGGTA
CCTCACAGAGAATATAACCGCTTCTCCCCAATCCAGCTGGAGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACA
TCATGCACAGCATCAATGGCTATGTTTTGTATAGTTTGCAGTTGTCAAGTTTGTGTCATGAGGTGGCATACTGGTACATT
CTAAGCTTTGGAGCAGACTGACTTCTTCTGTCTTCTTCTGATATACCTTCAAACACAAAATGGTCTATGAAGA
CACACTCACCTTATCCCAATTTCTAGGAGAACTGTCTTCAATGTGATGGAACCCAGGTCTATGGATTCTGGGGTGCC
ACAACCTCAGACTTTTGGAAACAGAGGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTGGTGATTATTAC
GAGGACAGTTATGAAGATATTTAGCATACTTGCTGAGTAAAAACAATGCCATTGAACCAAGAAGCTTCGAAATAACTCG
TACTACTCTTCAGTCAGATCAAGAGGAAATTTGACTATGATGATAACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACA
TTTATGATGAGGATGAAATCAGAGCCCCCGCAGCTTTCAAAGAAAACACGACACTATTTTATTGCTGCAGTGGAGAGG
CTCTGGGATTATGGGATGAGTAGCTCCCCACATGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAA
AGTTGTTTTCCAGGAATTTACTGATGGCTCCTTACTCAGCCCTTATACCGTGGAGAACTAAATGAACATTTGGGACTCC
TGGGGCCATATATAAGAGCAGAAGTTGAAGATAATATCATGGTAACCTTTAGAAATCAGGCCTCTCGTCCCTATTCCCTC
TATTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGAAAAAACTTTGTCAAGCCTAATGAAAC
CAAACTTACTTTTGGAAAGTGCAACATCATATGGCACCCACTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCT
CTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCTGGTCTGCCACACTAACACACTGAACCCCT
GCTCATGGGAGACAAGTGACAGTACAGGAATTTGCTCTGTTTTTACCATCTTTGATGAGACCAAAGCTGGTACTTCAC
TGAAATATGGAAGAACTGCAGGGCTCCCTGCAATATCCAGATGGAAGATCCCCTTTTAAAGAGAATTATCGCTTCC
ATGCAATCAATGGCTACATAATGGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTCGATGGTATCTGCTC
AGCATGGGCAGCAATGAAACATCCATTCTATTCAATTCAGTGACATGTGTTCACTGTACGAAAAAAGAGGAGTATAA
AATGGCACTGTACAATCTCTATCCAGGTGTTTTTGGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTGGCGGGTGG
AATGCCTTATTGGCGAGCATCTAGATGCTGGGATGAGCACACTTTTCTGGGTACAGCAATAAGTGTAGACTCCCCCTG
GGAATGGCTTCTGGACACATTAAGATTTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCAAAGCTGGCCAG
ACTTCATTATTCCGGATCAATCAATGCCTGGAGCACCAAGGAGCCCTTTTCTTGGATCAAGGTGGATCTGTTGGCACCAA
TGATTATTACGGCATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTTTATCATCATGTAT
AGTCTTGATGGGAAGAAGTGGCAGACTTATCGAGGAAATCCACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTCT
ATCTGGGATAAAACACAATATTTTTAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAACTCATTATAGCATTC

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

TGTATAAATTCTCTAAGAAACACGCATCTTTATAGAAACGTCCTATGATAGAGTTGAAATCAAGAGAAATCACATTTTCAGCAAT
 ACAGGGGAAAATCTTGCTAAAGCAGGAGTTTTCCGATGGGTACAAATATCCATGAACATAAAAGATATTACTATACCTTT
 GATAATTCACTACTATTTACTGAGAGCATTCAGAACACTACACAAATCTTTCCACGCTAAATCATAACGTCGGTTTCTT
 CCGTGTGACGACCGGGGCGTTGGCATAATGCAATACGTGTACGCGCTAAACCCTGTGTGCATCGTTTTAATTATTTCCCGG
 AACTCCCCGAGAGAAGTTCCCCGTCAGGGCTGTGGACATAGTTAATCCGGGAATACAATGACGATTTCATCGCACCTGAC
 ATACATTAATAAATATTAACAATATGAAATTTCAACTCATTGTTTATAGGTTTGTTTAATTTTCTACACATACGATTCTGTC
 GAACTTCAAAAAGCATCGGGAATAACACCATGAAAAAATGCTACTCGCTACTGCGCTGGCCCTGCTTATTACAGGATGT
 GCTCAACAGACGTTTACTGTTCAAAACAAACCGGCAGCAGTAGCACCAAAGGAAACCATCACCCATCATTTCTTCGTTTC
 TGGAAATTGGGCAGAAGAAAACCTGTCGATGCAGCCAAAATTTGTGGCGGCGCAGAAAATGTTTGTAAACAGAAACCCAGC
 AAACATTCGTAAATGGATTGCTCGGTTTTATTACTTTAGGCATTTATACTCCGCTGGAAGCGCGTGTGTATTGCTCACAA
 TAATTGCATGAGTTGCCCATCGCGATATGGGCAACTCTATCTGCACCTGCTCATTAAATATACTTCTGGGTTCCCTCCAGTT
 GTTTTTGTCATAGTGATCAGCCTCTCTCTGAGGGTGAAATAATCCCGTTTCAGCGGTGTCTGCCAGTCGGGGGGAGGCTGCA
 TTATCCACGCCGGAGGCGGTGGTGGCTTCACGCACTGACTGACAGACTGCTTTTGATGTGCAACCGACGACGACCAGCGGC
 AACATCATCACGCAGAGCATCATTTTCAGCTTTAGCATCAGCTAACTCCTTCGTGTATTTTGCATCGAGCGCAGCAACAT
 CACGCTGACGCATCTGCATGTCAGTAATTGCCGCGTTCGCCAGCTTCAGTTCTCTGGCATTTTTGTGCGCTGGGCTTTG
 TAGGTAATGGCGTTATCACGGTAATGATTAACAGCCCATGACAGGCAGACGATGATGCAGATAACCAGAGCGGAGATAAT
 CGCGGTGACTCTGCTCATAACATCAATCTCTCTGACCGTTCGGCCCGCTTCTTTGAATTTTGAATCAGGCTGTGAGCCTT
 ATGCTCGAACTGACCATAACAGCAGCCCGGCAGTGAAGCCAGATATTGCTGCAACCGTTCGATTGCTGACGGATATCAC
 CAGCATCAACTCATAGGTAAGCGCCACGCTCCTTTAATCTGCTGCAATGCCACAGCGTCTCGATTCTTCGGAGAGAAGTCT
 TTCAGGCCAAGCTGCTTGCGGTAGGCATCCCAACCGGAAAGAAGCTGGTAGCGTCCGGCGCCTGTTGATTGTAGTTT
 TGGGTTTTAGCGTGACAAGTTTGCAGGGGTGATCGGAGTAATCAGTAAATAGCTCTCCGCCCTACAATGACGTCATAACCAT
 GATTTCTGGTTTTTCTGACGTCCGTTATCAGTTCCCTCCGACCACGCCAGCATATCGAGGAACGCCTTACGTTGATTATTG
 ATTTCTACCATCTTCTACTCCGGCTTTTTTTAGCAGCGAAGCGTTTTGATAAGCGAACCAATCGAGTCAGTACCGATGTAGC
 CGATAAACACGCTCGTTATATAAGCGAGATTGCTACTTTAGTCCGGCGAAGTCGAGAAGGTCACGAATGAACCAGGCGATA
 ATGGCGCACATCGTTGCGTCGATTACTGTTTTTTGTAAACGCACCGCCATTATATCTGCCGCGAAGGTACGCCATTGCAAA
 CGCAAGGATTGCCCCGATGCCTTGTTCTTTGCCGCGAGAATGGCGGCCAACAGGTCATGTTTTTCTGGCATCTTCATGT
 CTTACCCCCAATAAGGGGATTGTCTCTATTTAATTAGGAATAAGGTCGATTACTGATAGAACAAATCCAGGCTACTGTGT
 TTAGTAATCAGATTGTTCGTGACCGATATGCACGGGCAAACGGCAGGAGGTTGTTAGCGCGACCTCCTGCCACCCGCT
 TTCACGAAGTCATGTGTAAGGCGCGCAGCGTAACTATTACTAATGAATTGAGGACAGACAGTGGCTACGGCTCAGTTT
 GGGTTGTGCTGTTGCTGGGCGGCGATGACGCCTGTACGCATTTGGTGATCCGGTTCCTGCTTCCGGTATTTCGCTTAATTCA
 GCACAACGGAAAGAGCACTGGCTAACCAGGCTCGCCGACTCTTCACGATTATCGACTCAATGCTCTTACCTGTTGTGCAG
 ATATAAAAAATCCCCGAAACCGTTATGCAGGCTCTAACTATTACCTGCGAACTGTTTCGGGATTGCATTTTGCAGACCTCT
 CTGCCTGCGATGGTTGGAGTTCCAGACGATACGTCGAAGTGACCAACTAGGCGGAATCGGTAGTAAGCGCCGCTCTTTT
 CATCTCACTACCACAACGAGCGAATTAACCCATCGTTGAGTCAAATTTACCCAATTTTATTCAATAAGTCAATATCATGC
 CGTTAATATGTTGCCATCCGTGGCAATCATGCTGCTAACGTGTGACCGCATTCAAATGTTGCTGCGATTGACTCTTCT
 TTGTGGCATTGCACCACCAGAGCGTCATACAGCGGCTTAACAGTGCGTGACCGAGGTGGGTTGGGTAAGGTTTGGGATTAG
 CATCGTCACAGCGGATATGCTGCGCTTGCTGGCATCCTTGAATAGCCGACGCCTTTCATCTTCCGCACTCTTTCTCGA
 CAACTCTCCCCACAGCTCTGTTTTGGCAATATCAACCGCACGGCCTGTACCATGGCAATCTCTGCATCTTGCCCCCGGC
 CTCGCGGCATACGGCAATAATCCGCATAAGCGAATGTTGCGAGCACTTGCAGTACCTTTGCCTTAGTATTTCCTTCAAG
 CTGCCCCCTGCAGG

FIGURE 6

CGCCCCCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAA
AGCCCCGGGCGTCGGGCGACCTTTGGTTCGCCCGGCCTCAGTGAGCGAGCGAGC
GCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGCCGCACG
CGTGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCT
TTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAAC
GTTCTTTTTTCGCAACGGGTTTGCCGCCCGCGGCAGGTAAGTGCCAGGGAAT
GTTTGTTCTTAAATACCATCGCTCCAGGGAATGTTTGTTCTTAAATACCATCT
ACTGACACTGACATCCACTTTTTCTTTTTCTCCACAGGTATCGATCCACCATGC
AAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCGATTCTGCTTTAGTG
CCACCAGAAGATACTACCTGGGTGCAGTGGAAGTGTATGGGACTATATGCA
AAGTGATCTCGGTGAGCTGCCTGTGGACGCAAGATTTCTCCTAGAGTGCCA
AAATCTTTTCCATTCAACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAGA
ATTCACGGATCACCTTTTCAACATCGCTAAGCCAAGGCCACCCTGGATGGGTC
TGCTAGGTCCTACCATCCAGGCTGAGGTTTATGATACAGTGGTCATTACACTT
AAGAACATGGCTTCCCATCCTGTCAGTCTTCATGCTGTTGGTGTATCCTACTG
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAA
AGAAGATGATAAAGTCTTCCCTGGTGGAAGCCATACATATGTCTGGCAGGTC
CTGAAAGAGAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATA
TCTTTCTCATGTGGACCTGGTAAAAGACTTGAATTCAGGCCTCATTGGAGCCC
TACTAGTATGTAGAGAAGGGAGTCTGGCCAAGGAAAAGACACAGACCTTGCA
CAAATTTATACTACTTTTTTGCTGTATTTGATGAAGGGAAAAGTTGGCACTCAG
AAACAAAGAACTCCTTGATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTG
GCCTAAAATGCACACAGTCAATGGTTATGTAAACAGGTCTCTGCCAGGTCTG
ATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTGGAATGGGCACCA
CTCCTGAAGTGCCTCAATATTCCTCGAAGGTCACACATTTCTTGTGAGGAAC
CATCGCCAGGCGTCCTTGGAATCTCGCCAATAACTTTCCTTACTGCTCAAAC
ACTCTTGATGGACCTTGGACAGTTTCTACTGTTTTGTATATCTCTTCCCACCA
ACATGATGGCATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCC
CAACTACGAATGAAAAATAATGAAGAAGCGGAAGACTATGATGATGATCTTA
CTGATTCTGAAATGGATGTGGTCAGGTTTGATGATGACAACTCTCCTTCCTTT
ATCCAAATTCGCTCAGTTGCCAAGAAGCATCCTAAAACCTTGGGTACATTACAT
TGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTCCTCGCCCCGATG
ACAGAAGTTATAAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAG
GAAGTACAAAAAAGTCCGATTTATGGCATAACACAGATGAAACCTTTAAGACT
CGTGAAGCTATTCAGCATGAATCAGGAATCTTGGGACCTTTACTTTATGGGG
AAGTTGGAGACACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATA
TAACATCTACCTCACGGAATCACTGATGTCCGTCCTTTGTATTCAAGGAGAT
TACCAAAGGTGTAAAACATTTGAAGGATTTTCCAATTCTGCCAGGAGAAAT
ATTCAAATATAAATGGACAGTGACTGTAGAAGATGGGCCAACTAAATCAGAT
CCTCGGTGCCTGACCCGCTATTACTCTAGTTTCGTTAATATGGAGAGAGATCT
AGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTAGATC
AAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGT
ATTTGATGAGAACCGAAGCTGGTACCTCACAGAGAATATACAACGCTTTCTC

CCCAATCCAGCTGGAGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACA
TCATGCACAGCATCAATGGCTATGTTTTTGATAGTTTGCAGTTGTCAGTTTGT
TTGCATGAGGTGGCATACTGGTACATTCTAAGCATTGGAGCACAGACTGACT
TCCTTTCTGTCTTCTTCTCTGGATATACCTTCAAACACAAAATGGTCTATGAA
GACACACTCACCTATTCCCATTCTCAGGAGAAACTGTCTTCATGTGCGATGGA
AAACCCAGGTCTATGGATTCTGGGGTGCCACAACTCAGACTTTCGGAACAGA
GGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTGGTGATT
ATTACGAGGACAGTTATGAAGATATTTTCAGCATACTTGCTGAGTAAAAACAA
TGCCATTGAACCAAGAAGCTTCTCCCAGAATCCACCAGTCTTGAAACGCCATC
AACGCGAAATAACTCGTACTACTCTTCAGTCAGATCAAGAGGAAATTGACTA
TGATGATACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACATTTATGAT
GAGGATGAAAATCAGAGCCCCCGCAGCTTTCAAAAGAAAACACGACACTATT
TTATTGCTGCAGTGGAGAGGCTCTGGGATTATGGGATGAGTAGCTCCCCACA
TGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAAAGTT
GTTTTCCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGGAGA
ACTAAATGAACATTTGGGACTCCTGGGGCCATATATAAGAGCAGAAGTTGAA
GATAATATCATGGTAACTTTCAGAAATCAGGCCTCTCGTCCCTATTCCTTCTA
TTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGA
AAAAACTTTGTCAAGCCTAATGAAACCAAACTTACTTTTGGAAGGTGCAAC
ATCATATGGCACCCACTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTT
TCTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCT
GGTCTGCCACACTAACACACTGAACCCTGCTCATGGGAGACAAGTGACAGTA
CAGGAATTTGCTCTGTTTTTACCATCTTTGATGAGACCAAAGCTGGTACTT
CACTGAAAATATGGAAAGAACTGCAGGGCTCCCTGCAATATCCAGATGGAA
GATCCCACTTTTAAAGAGAATTATCGCTTCCATGCAATCAATGGCTACATAAT
GGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTCGATGGTAT
CTGCTCAGCATGGGCAGCAATGAAAACATCCATTCTATTTCATTTAGTGGACA
TGTGTTCACTGTACGAAAAAAGAGGAGTATAAAATGGCACTGTACAATCTC
TATCCAGGTGTTTTTGAGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTG
GCGGGTGGAATGCCTTATTGGCGAGCATCTACATGCTGGGATGAGCACACTT
TTTCTGGTGTACAGCAATAAGTGTGCACTCCCCTGGGAATGGCTTCTGGACA
CATTAGAGATTTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCA
AAGCTGGCCAGACTTCATTATTCCGGATCAATCAATGCCTGGAGCACCAAGG
AGCCCTTTTCTTGGATCAAGGTGGATCTGTTGGCACCAATGATTATTCACGGC
ATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTT
TATCATCATGTATAGTCTTGATGGGAAGAAGTGGCAGACTTATCGAGGAAAT
TCCACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTTCATCTGGGATAAA
ACACAATATTTTTTAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAA
CTCATTATAGCATTTCGCAGCACTCTTCGCATGGAGTTGATGGGCTGTGATTTA
AATAGTTGCAGCATGCCATTGGGAATGGAGAGTAAAGCAATATCAGATGCAC
AGATTACTGCTTCATCCTACTTTACCAATATGTTTGCCACCTGGTCTCCTTCAA
AAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTGGAGACCTCAGGTGAA
TAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAAGTCACA
GGAGTAACTACTCAGGGAGTAAAATCTCTGCTTACCAGCATGTATGTGAAGG
AGTTCCTCATCTCCAGCAGTCAAGATGGCCATCAGTGGACTCTTTTTTTTCAG
AATGGCAAAGTAAAGGTTTTTTCAGGGAAATCAAGACTCCTTCACACCTGTGG

TGA ACTCTCTAGACCCACCGTTACTGACTCGCTACCTTCGAATTCACCCCCAG
AGTTGGGTGCACCAGATTGCCCTGAGGATGGAGGTTCTGGGCTGCGAGGCAC
AGGACCTCTACTGACTCGAGCCTAATAAAGGAAATTTATTTTCATTGCAATAG
TGTGTTGGTTTTTTTGTGTGCGGCCGCAGGAACCCCTAGTGATGGAGTTGGCCA
CTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGACCAAAGGTCGCC
CGACGCCCCGGGCTTTGCCCGGGCGGCCTCAGTGAGCGAGCGAGCGCGCAGCT
GCCTGCAGGACAT

FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10





FIGURE 11



DECLARATION FOR PATENT APPLICATION

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated next to my name. I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled **ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS**, the specification of which is attached hereto. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

<u>09/364,862</u>	<u>07/30/99</u>	<u>Pending</u>
<i>Application Serial No.</i>	<i>Filing Date</i>	<i>Patented, Pending or Abandoned</i>
<u>60/125,974</u>	<u>03/24/99</u>	<u>Pending</u>
<i>Application Serial No.</i>	<i>Filing Date</i>	<i>Patented, Pending or Abandoned</i>
<u>60/104,994</u>	<u>10/20/98</u>	<u>Abandoned</u>
<i>Application Serial No.</i>	<i>Filing Date</i>	<i>Patented, Pending or Abandoned</i>

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under § 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of Sole or First Inventor: **Linda B. Couto**

Inventor's Signature: _____
Residence: 7814 Oak Creek Drive, Pleasanton, California 94588
Post Office Address: 7814 Oak Creek Drive, Pleasanton, California 94588

Date: _____
Citizenship: United States of America

Full Name of Second Joint Inventor: **Peter C. Colosi**

Inventor's Signature: _____
Residence: 2016 Encinal Avenue, Alameda, California 94501
Post Office Address: 2016 Encinal Avenue, Alameda, California 94501

Date: _____
Citizenship: United States of America

Full Name of Third Joint Inventor: **Xiaobing Qian**

Inventor's Signature: _____
Residence: 3310 Fir Avenue, Alameda, California 94502
Post Office Address: 3310 Fir Avenue, Alameda, California 94502

Date: _____
Citizenship: P.R. China

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Linda B. Couto *et al.* Group No.:
Serial No.: Examiner:
Filed: 12/22/99
Entitled: **ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS**

POWER OF ATTORNEY BY ASSIGNEE

Assistant Commissioner for Patents
Washington, D.C. 20231

Avigen, Inc., as Assignee of record of the entire interest of the above-identified patent application, hereby appoints the members of the firm of MEDLEN & CARROLL, LLP, a firm composed of:

Virginia S. Medlen	(Reg. No. 32,050)	Maha A. Hamdan	(Reg. No. 43,655)
Peter G. Carroll	(Reg. No. 32,837)	J. Mitchell Jones	(Reg. No. 44,174)
Kamrin T. MacKnight	(Reg. No. 38,230)	David J. Wilson	(Reg. No. 45,225)
David A. Casimir	(Reg. No. 42,395)	Jason R. Bond	(Reg. No. P45,439)

as its attorneys with full power of substitution to prosecute this application and transact all business in the Patent and Trademark Office in connection therewith.

Please direct all future correspondence and telephone calls regarding this application to:

Kamrin T. MacKnight	
MEDLEN & CARROLL, LLP	
220 Montgomery Street, Suite 2200	Telephone: 415/705-8410
San Francisco, California 94104	Facsimile: 415/397-8338

I hereby certify that the Assignment document filed with the application or filed subsequent to the filing date of the application, has been reviewed and I hereby certify that, to the best of my knowledge and belief, title is with Avigen, Inc.

Dated: _____

By: _____

Name: _____

Title: _____

Avigen, Inc.
1201 Harbor Bay Parkway, #1000
Alameda, California 94502

SEQUENCE LISTING

<110> Couto, Linda B.
 Colosi, Peter C.
 5
 <120> Adeno-Associated Vectors for Expression of Factor VIII
 by Target Cells
 <130> Avigen-04082
 10
 <140> xx/xxx,xxx
 <141> 1999-12-22
 <150> 09/364,862
 15 <151> 1999-07-30
 <150> 60/125,974
 <151> 1999-03-24
 20 <150> 60/104,994
 <151> 1998-10-20
 <160> 15
 25 <170> PatentIn Ver. 2.0
 <210> 1
 <211> 59
 <212> DNA
 30 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 35 <400> 1
 cccaagcttg cggccgccc ggtgccccc ctaggcaggt aagtgccgtg tgtgggtcc 59
 <210> 2
 <211> 59
 40 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 45 <400> 2
 ccgctcgagc agagctctat ttgcatggtg gaatcgatgc cgcggaacc acacacggc 59
 <210> 3
 50 <211> 103
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

<400> 3
 5 cccaagcttg cggccgcccc ggtgccgccc ctaggcaggt aagtgccgtg tgtgggtccc 60
 gcggcatcga ttccaccatg caaatagagc tctgctcgag cgg 103

<210> 4
 <211> 57
 10 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

15 <400> 4
 ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactga 57

<210> 5
 20 <211> 57
 <212> DNA
 <213> Artificial Sequence

<220>
 25 <223> Description of Artificial Sequence: Synthetic

<400> 5
 gaatcgatac ctgtggagaa aaagaaaaag tggatgtcag tgtcagtaat tcaaggc 57

30 <210> 6
 <211> 99
 <212> DNA
 <213> Artificial Sequence

35 <220>
 <223> Description of Artificial Sequence: Synthetic

<400> 6
 40 ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactgacac 60
 tgacatccac tttttctttt tctccacagg tatcgattc 99

<210> 7
 <211> 100
 <212> DNA
 45 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

50 <400> 7
 aggggaatggt tgtttttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60
 aatgtttggt cttaaatacc atctacagtt attggttaaa 100

<210> 8
 <211> 59
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 8
 10 ggaaaggtga tctgtgtgca gaaagactcg ctctaataa cttctttaac caataactg 59
 <210> 9
 <211> 144
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 9
 20 agggaatggt tgttcttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60
 aatgtttggt cttaaatacc atctacagtt attggttaaa gaagtatatt agagcgagtc 120
 tttctgcaca cagatcacct ttcc 144
 25 <210> 10
 <211> 59
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 10
 35 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgc 59
 <210> 11
 <211> 59
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <223> Description of Artificial Sequence: Synthetic
 <400> 11
 45 tcgagcggcc gcacacaaaa aaccaacaca cagatctcta gagctctgat cttttattc 59
 <210> 12
 <211> 63
 <212> DNA
 50 <213> Artificial Sequence
 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 12

5 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgct 60
cga 63

<210> 13

<211> 11933

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

15 <400> 13

cagctgcgcg ctgcctcgt cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc 60
tttggctgcc cggcctcagt ggcgcagcga gcgcgcagag agggagtggc caactccatc 120
actaggggtt cctgcggccg cccaggggaat gtttgttctt aaataccatc cagggaatgt 180
20 ttgttcttaa ataccatcca gggaatgttt gttcttaaata accatctaca gttattgggt 240
aaagaagtat attagagcga gtctttctgc acacagatca cctttccggg tgcgcgccct 300
aggcaggtaa gtgccgtgtg tggttccgc gccctggcc tctttacggg ttatggccct 360
tgcgtgcctt gaattactga cactgacatc cactttttct tttctccac aggtatcgat 420
tccaccatgc aaatagagct ctccacctgc ttctttctgt gccttttgcg attctgcttt 480
agtgccacca gaagatacta cctgggtgca gtggaactgt catgggacta tatgcaaagt 540
25 gatctcgggtg agctgcctgt ggacgcaaga tttcctccta gactgcaaaa atcttttcca 600
ttcaacacct cagtctgtga caaaaagact ctgtttgtag aattcacgga tcaccttttc 660
aacatcgcta agccaaggcc accctggatg ggtctgctag gtccatccat ccaggctgag 720
gtttatgata cagtggctcat tacacttaag aacatggctt cccatcctgt cagtcttcat 780
gctgttggtg tctctactg gaaagcttct gagggagctg aatatgatga tcagaccagt 840
30 caaagggaga aagaagatga taaagtcttc cctgggtgaa gccatacata tgtctggcag 900
gtcctgaaag agaattggtcc aatggcctct gaccactgt gccttaccta ctcatatctt 960
tctcatgtgg acctggtaaa agacttgaat tcaggcctca ttggagccct actagtatgt 1020
agagaaggga gtctggccaa ggaaaagaca cagaccttgc acaaatttat actacttttt 1080
gctgtatttg atgaaggaa aagttggcac tcagaaacaa agaactcctt gatgcaggat 1140
35 agggatgctg catctgctcg ggcctggcct aaaatgcaca cagtcaatgg ttatgtaaac 1200
aggctctctg caggctctgat tggatgccac aggaaatcag tctattggca tgtgattgga 1260
atgggcacca ctctgaagt gcactcaata ttctcgaag gtcacacatt tcttgtgagg 1320
aaccatcgcc aggcgtcctt ggaaatctcg ccaataactt tcttactgc tcaaactc 1380
ttgatggacc ttggacagtt tctactgttt tgtcatatct ctccaccacacatgatggc 1440
40 atggaagctt atgtcaaagt agacagctgt ccagaggaac cccaactacg aatgaaaaat 1500
aatgaagaag cggaagacta tgatgatgat ctactgatt ctgaaatgga tgtggctcagg 1560
tttgatgatg acaactctcc ttcttttctc caaattcgct cagttgccaa gaagcatcct 1620
aaaacttggg tacattacat tgctgctgaa gaggaggact gggactatgc tcccttagtc 1680
ctgcgcccg atgacagaag ttataaaagt caatatttga acaatggccc tcagcggatt 1740
45 ggtaggaagt acaaaaaagt ccgatttatg gcatacacag atgaaacctt taagactcgt 1800
gaagctattc agcatgaatc aggaatcttg ggacctttac tttatgggga agttggagac 1860
aactgttga ttatatttaa gaatcaagca agcagaccat ataactcta cctcacgga 1920
atcactgatg tccgtccttt gtattcaagg agattaccaa aaggtgtaaa acatttgaag 1980
gattttccaa ttctgccagg agaaatattc aaatataaat ggacagtgc tgtagaagat 2040
50 gggccaacta aatcagatcc tcggtgcctg acccgctatt actctagttt cgtaaatatg 2100
gagagagatc tagcttcagg actcattggc cctctcctca tctgctacaa agaactctga 2160
gatcaaagag gaaaccagat aatgtcagac aagaggaatg tcatcctggt ttctgtattt 2220

	gatgagaacc	gaagctggta	cctcacagag	aatatacaac	gcttttctccc	caatccagct	2280
	ggagtgcagc	ttgaggatcc	agagttccaa	gcctccaaca	tcatgcacag	catcaatggc	2340
	tatgtttttg	atagtttgca	gttgtcagtt	tgtttgcatg	aggtggcata	ctggtacatt	2400
	ctaagcattg	gagcacagac	tgacttcctt	tctgtcttct	tctctggata	taccttcaaa	2460
5	cacaaaatgg	tctatgaaga	cacactcacc	ctattcccat	tctcaggaga	aactgtcttc	2520
	atgtcgatgg	aaaaccagg	tctatggatt	ctggggtgcc	acaactcaga	ctttcggaac	2580
	agaggcatga	ccgccttact	gaaggtttct	agttgtgaca	agaacactgg	tgattattac	2640
	gaggacagtt	atgaagatat	ttcagcatac	ttgctgagta	aaaacaatgc	cattgaacca	2700
	agaagcttcg	aaataactcg	tactactctt	cagtcagatc	aagaggaaat	tgactatgat	2760
10	gataccatat	cagttgaaat	gaagaaggaa	gattttgaca	tttatgatga	ggatgaaaat	2820
	cagagccccc	gcagctttca	aaagaaaaca	cgacactatt	ttattgctgc	agtggagagg	2880
	ctctgggatt	atgggatgag	tagctcccca	catgttctaa	gaaacagggc	tcagagtggc	2940
	agtgtccctc	agttcaagaa	agttgttttc	caggaattta	ctgatggctc	ctttactcag	3000
	cccttatacc	gtggagaact	aatgaacat	ttgggactcc	tgggggccata	tataagagca	3060
15	gaagttgaag	ataatatcat	ggtaactttc	agaaatcagg	cctctcgctc	ctattccttc	3120
	tattctagcc	ttatttctta	tgaggaagat	cagaggcaag	gagcagaacc	tagaaaaaac	3180
	tttgtcaagc	ctaataaacc	caaaacttac	ttttggaaag	tgcaacatca	tatggcaccc	3240
	actaaagatg	agtttgactg	caaagcctgg	gcttatttct	ctgatgttga	cctggaaaaa	3300
	gatgtgcact	caggcctgat	tggacccctt	ctggctctgc	acactaacac	actgaaccct	3360
20	gctcatggga	gacaagtgc	agtacaggaa	tttgctctgt	ttttcaccat	ctttgatgag	3420
	acaaaaagct	ggtacttcac	tgaaaatatg	gaaagaaact	gcagggctcc	ctgcaatatc	3480
	cagatggaag	atcccacttt	taaagagaat	tatcgcttcc	atgcaatcaa	tggtacata	3540
	atggatacac	tacctggctt	agtaatggct	caggatcaaa	ggattcgatg	gtatctgctc	3600
	agcatgggca	gcaatgaaaa	catccattct	attcatttca	gtggacatgt	gttcaactgta	3660
25	cgaaaaaaaag	aggagtataa	aatggcactg	tacaatctct	atccagggtg	ttttgagaca	3720
	gtggaaatgt	taccatccaa	agctggaatt	tggcggtggg	aatgccttat	tggcgagcat	3780
	ctacatgctg	ggatgagcac	actttttctg	gtgtacagca	ataagtgtca	gactcccctg	3840
	ggaatggctt	ctggacacat	tagagatttt	cagattacag	cttcaggaca	atatggacag	3900
	tgggccccaa	agctggccag	acttcattat	tccggatcaa	tcaatgcctg	gagcaccaag	3960
30	gagccctttt	cttggatcaa	ggtggatctg	ttggcaccaa	tgattattca	cggcatcaag	4020
	acccagggtg	cccgctcagaa	gttctccagc	ctctacatct	ctcagtttat	catcatgtat	4080
	agtcttgatg	ggaagaagtg	gcagacttat	cgaggaaatt	ccactggaac	cttaatggtc	4140
	ttctttggca	atgtggattc	atctgggata	aaacacaata	tttttaaccc	tccaattatt	4200
	gctcgataca	tccgtttgca	cccaactcat	tatagcattc	gcagcactct	tgcgatggag	4260
35	ttgatgggct	gtgatttaaa	tagttgcagc	atgccattgg	gaatggagag	taaagcaata	4320
	tcagatgcac	agattactgc	ttcatcctac	tttaccata	tgtttgccac	ctggtctcct	4380
	tcaaaagctc	gacttcacct	ccaagggagg	agtaatgcct	ggagacctca	ggtgaataat	4440
	ccaaaagagt	ggctgcaagt	ggacttccag	aagacaatga	aagtcacagg	agtaactact	4500
	cagggagtaa	aatctctgct	taccagcatg	tatgtgaagg	agttcctcat	ctccagcagt	4560
40	caagatggcc	atcagtggac	tctctttttt	cagaatggca	aagtaaagg	ttttcaggg	4620
	aatcaagact	ccttcacacc	tgtggtgaac	tctctagacc	caccgttact	gactcgctac	4680
	cttcgaattc	acccccagag	ttgggtgcac	cagattgcc	tgaggatgga	ggttctgggc	4740
	tgcgagggcac	aggacctcta	ctgactcgag	aataaaaagat	cagagctcta	gagatctgtg	4800
	tggtgggtttt	ttgtgtgcgg	ccgcaggaac	ccctagtgat	ggagtgggcc	actccctctc	4860
45	tgcgcgctcg	ctcgctcact	gaggccgggc	gaccaaagg	cgcccgacgc	ccgggctttg	4920
	cccgggcggc	ctcagtgagc	gagcgagcgc	gcagctgcct	gcaggacatg	tgagcaaaa	4980
	gccagcaaaa	ggccaggaac	cgtaaaaagg	ccgcgttgct	ggcgtttttc	cataggctcc	5040
	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	gaggtggcga	aacccgacag	5100
	gactataaag	ataccaggcg	tttccccctg	gaagctccct	cgtgcgctct	cctgttccga	5160
50	ccctgccgct	taccggatac	ctgtccgctt	ttctcccttc	gggaagcgtg	gcgctttctc	5220
	atagctcacg	ctgtaggtat	ctcagttcgg	tgtaggtcgt	tcgctccaag	ctgggctgtg	5280
	tgcacgaacc	ccccgttcag	cccgaccgct	gcgccttatc	cggtaactat	cgtcttgagt	5340

	ccaacccggt	aagacacgac	ttatcgccac	tggcagcagc	cactggtaac	aggatttagca	5400
	gagcgaggta	tgtaggcggt	gctacagagt	tcttgaagt	gtggcctaac	tacggctaca	5460
	ctagaaggac	agtatttggg	atctgcgctc	tgtgaagcc	agttaccttc	ggaaaaagag	5520
	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cgggtggtttt	tttgtttgca	5580
5	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcttttgatc	ttttctacgg	5640
	ggctctgacgc	tcagtggaa	gaaaactcac	gttaagggat	tttgggtcatg	agattatcaa	5700
	aaaggatctt	cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca	atctaaagta	5760
	tatatgagta	aacttggtct	gacagttacc	aatgcttaat	cagtgaaggca	cctatctcag	5820
	cgatctgtct	atttcgttca	tccatagttg	cctgactccc	cgctcgtgtag	ataactacga	5880
10	tacggggagg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	5940
	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	agaagtggtc	6000
	ctgcaacttt	atccgcctcc	atccagctca	ttaattgttg	ccgggaagct	agagtaagta	6060
	gttcgccagt	taatagtttg	cgcaacgttg	ttgccattgc	tacaggcatc	gtggtgtcac	6120
	gctcgtcggt	tggtatggct	tcattcagct	ccggttccca	acgatcaagg	cgagttacat	6180
15	gatcccccat	gttgtgcaaa	aaagcgggta	gctccttcgg	tcttccgatc	gttgtcagaa	6240
	gtaagttggc	cgcagtggtta	tcactcatgg	ttatggcagc	actgcataat	tctcttactg	6300
	tcatgccatc	cgtaagatgc	ttttctgtga	ctggtgagta	ctcaaccaag	tcattctgag	6360
	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgtc	aatacgggat	aataccgcgc	6420
	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	cgaaaactct	6480
20	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	6540
	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	aggcaaaatg	6600
	ccgcaaaaaa	gggaataagg	gcgacacgga	aatgttgaat	actcatactc	ttcctttttc	6660
	aatattattg	aagcatttat	caggggttatt	gtctcatgag	cggatacata	tttgaatgta	6720
	tttagaaaaa	taaacaaata	gggggttcgc	gcacatttcc	ccgaaaagt	ccacctgacg	6780
25	tctaagaaac	cattattatc	atgacattaa	cctataaaaa	taggcgtatc	acgaggccct	6840
	ttcgtctcgc	gcgtttcggg	gatgacgggtg	aaaacctctg	acacatgcag	ctcccggaga	6900
	cggtcacagc	ttgtctgtaa	gcggatgccg	ggagcagaca	agcccgtcag	ggcgcgtcag	6960
	cgggtgttgg	cgggtgtcgg	ggctggctta	actatgcggc	atcagagcag	attgtactga	7020
	gagtgcacca	taaaattgta	aacgttaata	ttttgttaaa	attcgcgtta	aatttttgtt	7080
30	aaatcagctc	attttttaac	caataggccg	aaatcggcaa	aatcccttat	aatcaaaaag	7140
	aatagcccga	gatagggttg	agtgtgttgc	cagtttgga	caagagtcca	ctattaaaga	7200
	acgtggactc	caacgtcaaa	gggcgaaaaa	ccgtctatca	gggcgatggc	ccactacgtg	7260
	aaccatcacc	caaatcaagt	tttttggggg	cgaggtgccg	taaagcacta	aatcgggaacc	7320
	ctaaagggag	ccccgattt	agagcttgac	ggggaaagcc	ggcgaacgtg	gcgagaaagg	7380
35	aagggaaagaa	agcgaaagga	gcgggcgcta	gggcgctggc	aagtgtagcg	gtcacgctgc	7440
	gcgtaaccac	cacaccgcgc	gcgtttaatg	cgccgctaca	gggcgcgtac	tatggttgct	7500
	ttgacgtatg	cgggtgtgaa	taccgcacag	atgcgtaagg	agaaaatacc	gcacaggcc	7560
	gtaacctgtc	ggatcaccgg	aaaggaccgg	taaagtata	atgattatca	tctacatata	7620
	acaacgtgcg	tggaggccat	caaaccacgt	caaataatca	attatgacgc	aggtatcgta	7680
40	ttaattgata	tgcatcaact	taacgtaaaa	acaacttcag	acaatacaaa	tcagcgacac	7740
	tgaatacggg	gcaacctcat	gtcaacgaag	aacagaaccc	gcagaacaac	aaccgcgaac	7800
	atccgctttc	ctaaccaaat	gattgaacaa	attaacatcg	ctcttgagca	aaaaggggtcc	7860
	gggaattttc	cagcctgggt	cattgaagcc	tgccgtcgga	gactaacgtc	agaaaagaga	7920
	gcatatacat	caattaaaag	tgatgaagaa	tgaacatccc	gcgttcttcc	ctccgaacag	7980
45	gacgatattg	taaattcact	taattacgag	ggcattgcag	taattgagtt	gcagttttac	8040
	cactttcctg	acagtgcacg	actgcgtgtt	ggctctgtca	cagactaaat	agtttgaatg	8100
	attagcagtt	atggtgatca	gtcaaccacc	agggataaat	ccttcataatt	attatcgtgc	8160
	ttcaccaacg	ctgcctcaat	tgctctgaat	gcttcacagag	acaccttatg	ttctatacat	8220
	gcaattacaa	catcagggtta	actcatagaa	atggtgctat	taagcatatt	ttttacacga	8280
50	atcagatcca	cggagggatc	atcagcagat	tgttctttat	tcattttgtc	gctccatgcg	8340
	cttgctcttc	atctagcggg	taaaatatta	cttcaaattct	ttctgtatga	agatttgagc	8400
	acgttggcct	tacatacatc	tgtcgggtgt	atttccctcc	agaatgccag	caggaccgca	8460

	ctttgttacg	caaccaatac	tattaagtga	aaacattcct	aatatttgac	ataaatcatc	8520
	aacaaaacac	aaggaggtca	gaccagattg	aaacgataaa	aacgataatg	caaactacgc	8580
	gccctcgtat	cacatggaag	gtttttaccaa	tgggtcaggt	tgccattttt	aaagaaatat	8640
	tcgatcaagt	gcgaaaagat	ttagactgtg	aattgtttta	ttctgaacta	aaacgtcaca	8700
5	acgtctcaca	ttatatattac	tatctagcca	cagataatat	tcacatcgtg	ttagaaaacg	8760
	ataacaccgt	gttaataaaaa	ggacttaaaa	aggttgtaaa	tgttaaattc	tcaagaaaca	8820
	cgcattcttat	agaaacgtcc	tatgataggt	tgaaatcaag	agaaatcaca	tttcagcaat	8880
	acagggaaaa	tcttgctaaa	gcaggagttt	tccgatgggt	tacaaatatc	catgaacata	8940
	aaagatatta	ctataccttt	gataattcat	tactatttac	tgagagcatt	cagaacacta	9000
10	cacaaatctt	tcacgcctaa	atcataacgt	ccggtttctt	ccgtgtcagc	accggggcgt	9060
	tggcataatg	caatacgtgt	acgcgcctaaa	ccctgtgtgc	atcgttttta	ttattcccgg	9120
	acactcccgc	agagaagttc	cccgtcaggg	ctgtggacat	agttaatccg	ggaatacaat	9180
	gacgattcat	cgcacctgac	atacattaat	aaatattaac	aatatgaaat	ttcaactcat	9240
	tgtttagggg	ttgttttaatt	ttctacacat	acgattctgc	gaacttcaaa	aagcatcggg	9300
15	aataacacca	tgaaaaaaat	gctactcgct	actgcgctgg	ccctgcttat	tacaggatgt	9360
	gctcaacaga	cgtttactgt	tcaaaaacaaa	ccggcagcag	tagcaccaaa	ggaaaccatc	9420
	acccatcatt	tcttcgtttc	tgggaattggg	cagaagaaaa	ctgtcgatgc	agccaaaatt	9480
	tgtggcggcg	cagaaaatgt	tgttaaaaca	gaaaccagc	aaacattcgt	aatggattg	9540
	ctcggtttta	ttacttttagg	cattttatact	ccgttggaag	cgcgtgtgta	ttgctcacia	9600
20	taattgcatg	agttgcccac	cgcgatatgg	gcaactctat	ctgcaactgt	cattaatata	9660
	cttctgggtt	ccttcacggt	gtttttgcat	agtgatcagc	ctctctctga	gggtgaaata	9720
	atcccgttca	gcggtgtctg	ccagtcgggg	ggaggctgca	ttatccacgc	cggaggcggg	9780
	ggtggcttca	cgcactgact	gacagactgc	tttgatgtgc	aaccgacgac	gaccagcggc	9840
	aacatcatca	cgcagagcat	cattttcagc	tttagcatca	gctaactcct	tcgtgtattt	9900
25	tgcacgagc	gcagcaacat	cacgctgacg	catctgcatg	tcagtaattg	ccgcgttcgc	9960
	cagcttcagt	tctctggcat	ttttgtcgcg	ctgggctttg	taggtaattg	cgttatcacg	10020
	gtaatgatta	acagcccacg	acaggcagac	gatgatgcag	ataaccagag	cggagataat	10080
	cgcgggtgact	ctgctcatac	atcaatctct	ctgaccgttc	cgcccgttcc	tttgaatttt	10140
	gcaatcaggc	tgtcagcctt	atgctcgaac	tgaccataac	cagcgcgccg	cagtgaagcc	10200
30	cagatattgc	tgcaacggtc	gattgcttga	cggatatcac	cacgatcaat	cataggtaaa	10260
	gcgccacgct	ccttaactctg	ctgcaatgcc	acagcgtcct	gacttttcgg	agagaagtct	10320
	ttcaggccaa	gctgcttgcg	gtaggcatcc	caccaacggg	aaagaagctg	gtagcgtccg	10380
	gcgcctggtg	atlttgagttt	tgggttttagc	gtgacaagtt	tgcgaggggtg	atcggagtaa	10440
	tcagtaaaata	gctctccgccc	tacaatgacg	tcataaccat	gatttctggt	tttctgacgt	10500
35	ccgttatcag	ttccctccga	ccacgccagc	atatcgagga	acgccttacg	ttgattattg	10560
	atttctacca	tcttctactc	cggctttttt	agcagcgaa	cgtttgataa	gcgaaccaat	10620
	cgagtcahta	ccgatgtagc	cgataaacac	gctcgttata	taagcgagat	tgctacttag	10680
	tccggcgaa	tcgagaaggt	cacgaatgaa	ccaggcgata	atggcgacac	tcgttgcgtc	10740
	gattactgtt	tttgtaaacg	caccgccatt	atatctgccc	cgaaggtagc	ccattgcaaa	10800
40	cgcaaggatt	gccccgatgc	cttggttcctt	tgccgcgaga	atggcggcca	acaggatcatg	10860
	tttttctggc	atcttcatgt	cttaccoccca	ataaggggat	ttgctctatt	taattaggaa	10920
	taaggctgat	tactgataga	acaaatccag	gctactgtgt	ttagtaataca	gatttggttcg	10980
	tgaccgatat	gcacgggcaa	aacggcagga	ggttgtagc	gcgacctcct	gccacccgct	11040
	ttcacgaagg	tcatgtgtaa	aaggccgcag	cgtaactatt	actaatgaat	tcaggacaga	11100
45	cagtggctac	ggctcagttt	gggttggtgt	gttgctgggc	ggcgatgacg	cctgtacgca	11160
	tttggtgatc	cggttctgct	tccggtattc	gcttaattca	gcacaacgga	aagagcactg	11220
	gctaaccagg	ctcgccgact	cttcacgatt	atcgactcaa	tgctcttacc	tggttgtagc	11280
	atataaaaaa	ttccgaaacc	gttatgcagg	ctctaactat	tacctgcgaa	ctgtttcggg	11340
	attgcatttt	gcagacctct	ctgcctgcga	tggttgaggt	tcacagacgat	acgtcgaagt	11400
50	gaccaactag	gcggaatcgg	tagtaagcgc	cgctctttt	catctcacta	ccacaacgag	11460
	cgaattaacc	catcgttgag	tcaaattttac	ccaattttat	tcaataagtc	aatatcatgc	11520
	cgttaatatg	ttgccatccg	tggcaatcat	gctgctaacc	tgtagaccgca	ttcaaaatgt	11580

5 tgtctgcgat tgactcttct ttgtggcatt gcaccaccag agcgtcatac agcggcttaa 11640
 cagtgcgtga ccaggtgggt tgggtaaggt ttgggattag catcgtcaca gcgcgatatg 11700
 ctgcgcttgc tggcatcctt gaatagccga cgcctttgca tcttccgcac tctttctcga 11760
 caactctccc ccacagctct gttttggcaa tatcaaccgc acggcctgta ccatggcaat 11820
 ctctgcatct tgcccccggt gtcgcggcac tacggcaata atccgcataa gcgaatgttg 11880
 cgagcacttg cagtaccttt gccttagtat ttccttcaag ctgccccctgc agg 11933

<210> 14

<211> 4999

10 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

15

<400> 14

cgccccctgca ggcagctgct cgctcgctcg ctactgagg ccgccccggc aaagccccggg 60
 cgtcggggcga ccttttggtcg cccggcctca gtgagcgagc gagcgcgag agagggagtg 120
 gccaactcca tcaactagggg ttctcgcggc cgcacgcgtg gtggcgcggg gtaaaactggg 180
 20 aaagtgatgt cgtgtactgg ctccgccttt ttcccgaggg tgggggagaa ccgtatataa 240
 gtgcagtagt cgccgtgaac gttctttttc gcaacgggtt tgccgccccg cggcaggtaa 300
 gtgccaggga atgtttgttc ttaaatacca tcgctccagg gaatgtttgt tcttaaatac 360
 catctactga cactgacatc cactttttct tttctccac aggtatcgat ccaccatgca 420
 aatagagctc tccacctgct tctttctgtg ccttttgca ttctgcttta gtgccaccag 480
 25 aagatactac ctgggtgcag tggaaactgtc atgggactat atgcaaagt atctcgggtga 540
 gctgcctgtg gacgcaagat ttctcctag agtgccaaaa tcttttccat tcaacacctc 600
 agtcgtgtac aaaaagactc tgtttgtaga attcacggat caccttttca acatcgctaa 660
 gccaaggcca cctgggatgg gtctgctagg tcctaccatc caggctgagg tttatgatac 720
 agtggtcatt acacttaaga acatggcttc ccactctgtc agtcttcatg ctggttggtgt 780
 30 atcctactgg aaagcttctg agggagctga atatgatgat cagaccagtc aaagggagaa 840
 agaagatgat aaagtcttcc ctggtggaag ccatacatat gtctggcagg toctgaaaga 900
 gaatggtcca atggcctctg acccactgtg ccttacctac tcatatcttt ctcatgtgga 960
 cctggtaaaa gacttgaatt caggcctcat tggagcccta ctagtatgta gagaagggag 1020
 tctggccaag gaaaagacac agaccttgca caaatttata ctacttttg ctgtatttga 1080
 35 tgaagggaaa agttggcact cagaaacaaa gaactccttg atgcaggata gggatgctgc 1140
 atctgctcgg gcctggccta aaatgcacac agtcaatgggt tatgtaaaaca ggtctctgcc 1200
 aggtctgatt ggatgccaca ggaaatcagt ctattggcat gtgattggaa tgggcaccac 1260
 toctgaagtg cactcaatat tctcgaagg tcacacattt cttgtgagga accatcgcca 1320
 ggcgctcctt gaaatctcgc caataacttt cttactgct caaacactct tgatggacct 1380
 40 tggacagttt ctactgtttt gtcatatctc ttcccaccaa catgatggca tggaaagctta 1440
 tgtcaaagta gacagctgtc cagaggaacc ccaactacga atgaaaaata atgaagaagc 1500
 ggaagactat gatgatgata ttactgattc tgaaatggat gtggtcaggt ttgatgatga 1560
 caactctcct toctttatcc aaattcgctc agttgccaa aagcatccta aaacttgggt 1620
 acattacatt gctgctgaag aggaggactg ggactatgct cccttagtcc tcgcccccca 1680
 45 tgacagaagt tataaaagtc aatatttgaa caatggccct cagcggattg gtaggaagta 1740
 caaaaaagtc cgatttatgg catacacaga tgaaaccttt aagactcgtg aagctattca 1800
 gcatgaatca ggaatcttgg gacctttact ttatggggaa gttggagaca cactgttgat 1860
 tatatttaag aatcaagcaa gcagaccata taacatctac cctcacggaa tcaactgatg 1920
 ccgtcctttg tattcaagga gattacaaa aggtgtaaaa catttgaagg attttccaat 1980
 50 tctgccagga gaaatattca aatataaatg gacagtgact gtagaagatg ggccaactaa 2040
 atcagatcct cggtgctga cccgctatta ctctagtttc gttaatatgg agagagatct 2100
 agcttcagga ctcatgggcc ctctcctcat ctgctacaaa gaatctgtag atcaaagagg 2160

	aaaccagata	atgtcagaca	agaggaatgt	catcctgttt	tctgtatttg	atgagaaccg	2220
	aagctggtac	ctcacagaga	atatacaacg	ctttctcccc	aatccagctg	gagtgagct	2280
	tgaggatcca	gagttccaag	cctccaacat	catgcacagc	atcaatggct	atgtttttga	2340
	tagtttgag	ttgtcagttt	gtttgcatga	ggtagcatat	tggtacattc	taagcattgg	2400
5	agcacagact	gacttccttt	ctgtcttctt	ctctggatat	accttcaaac	acaaaatggg	2460
	ctatgaagac	acactcaccc	tattccatt	ctcaggagaa	actgtcttca	tgtcgatgga	2520
	aaaccaggt	ctatggattc	tggggtgcca	caactcagac	tttcggaaca	gaggcatgac	2580
	cgccttactg	aagggtttcta	gttgtgacaa	gaactcaggt	gattattacg	aggacagtta	2640
	tgaagatatt	tcagcatact	tgctgagtaa	aaacaatgcc	attgaaccaa	gaagcttctc	2700
10	ccagaatcca	ccagtcttga	aacgccatca	acgcgaaata	actcgtacta	ctcttcagtc	2760
	agatcaagag	gaaattgact	atgatgatac	catatcagtt	gaaatgaaga	aggaagattt	2820
	tgacatttat	gatgaggatg	aaaatcagag	ccccgcagc	tttcaaaaga	aaacacgaca	2880
	ctattttatt	gctgcagtg	agaggctctg	ggattatggg	atgagtagct	ccccacatgt	2940
	tctaagaaac	agggctcaga	gtggcagtg	ccctcagttc	aagaaagttg	ttttccagga	3000
15	atttactgat	ggctccttta	ctcagccctt	ataccgtgga	gaactaaatg	aacatttggg	3060
	actcctgggg	ccatatataa	gagcagaagt	tgaagataat	atcatggtta	ctttcagaaa	3120
	tcaggcctct	cgtccctatt	ccttctattc	tagccttatt	tcttatgagg	aagatcagag	3180
	gcaaggagca	gaacctagaa	aaaactttgt	caagcctaata	gaaacccaaa	cttacttttg	3240
	gaaagtgcaa	catcatatgg	caccactaa	agatgagttt	gactgcaaag	cctgggctta	3300
20	tttctctgat	gttgacctgg	aaaaagatgt	gcactcaggc	ctgattggac	cccttctggg	3360
	ctgccacact	aacacactga	accctgctca	tgggagacaa	gtgacagtac	aggaatttgc	3420
	tctgtttttc	accatctttg	atgagaccaa	aagctggtac	ttcactgaaa	atatggaaaag	3480
	aaactgcagg	gctccctgca	atatccagat	ggaagatccc	acttttaaaag	agaattatcg	3540
	cttccatgca	atcaatggct	acataatgga	tacactacct	ggcttagtaa	tggctcagga	3600
25	tcaaaggatt	cgatggtatc	tgctcagcat	gggcagcaat	gaaaacatcc	attctattca	3660
	tttcagtggg	catgtgttca	ctgtacgaaa	aaaagaggag	tataaaatgg	cactgtacaa	3720
	tctctatcca	ggtgtttttg	agacagtggg	aatgttacca	tccaaagctg	gaatttggcg	3780
	ggtggaatgc	cttattggcg	agcatctaca	tgctgggatg	agcacacttt	ttctgggtga	3840
	cagcaataag	tgtcagactc	ccctgggaat	ggcttctgga	cacattagag	attttcagat	3900
30	tacagcttca	ggacaatatg	gacagtgggc	cccaaagctg	gccagacttc	attattccgg	3960
	atcaatcaat	gcctggagca	ccaaggagcc	cttttcttgg	atcaaggtgg	atctgttggc	4020
	accaatgatt	attcacggca	tcaagaccca	gggtgcccg	cagaagttct	ccagcctcta	4080
	catctctcag	tttatcatca	tgtatagtct	tgatgggaag	aagtggcaga	cttatcgagg	4140
	aaattccact	ggaaccttaa	tggcttctct	tggcaatgtg	gattcatctg	ggataaaaca	4200
35	caatattttt	aacctccaa	ttattgctcg	atacatccgt	ttgcacccaa	ctcattatag	4260
	cattcgcagc	actcttcgca	tggagttgat	gggctgtgat	ttaaatagtt	gcagcatgcc	4320
	attgggaatg	gagagtaaag	caatatcaga	tgcacagatt	actgcttcat	cctactttac	4380
	caatatgttt	gccacctggg	ctccttcaaa	agctcgactt	cacctccaag	ggaggagtaa	4440
	tgcctggaga	cctcaggtga	ataatccaaa	agagtggctg	caagtggact	tccagaagac	4500
40	aatgaaagtc	acaggagtaa	ctactcaggg	agtaaaatct	ctgcttacca	gcatgtatgt	4560
	gaaggagttc	ctcatctcca	gcagtcaaga	tggccatcag	tggactctct	tttttcagaa	4620
	tggcaaagta	aagggttttc	agggaaatca	agactccttc	acacctgtgg	tgaactctct	4680
	agaccacccg	ttactgactc	gctaccttcg	aattcaccoc	cagagttggg	tgcaccagat	4740
	tgccttgagg	atggaggttc	tgggctgcga	ggcacaggac	ctctactgac	tcgagcctaa	4800
45	taaaggaaat	ttattttcat	tgcaatagtg	tgttggtttt	ttgtgtgcgg	ccgcagggaac	4860
	ccctagtgat	ggagttggcc	actccctctc	tgcgcgctcg	ctcgcctcact	gaggccgggc	4920
	gaccaaaggt	cgcgcgacgc	ccgggctttg	ccgggcgggc	ctcagtgagc	gagcgagcgc	4980
	gcagctgect	gcaggacat					4999
50	<210>	15					
	<211>	14					
	<212>	PRT					

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

5

<400> 15

Ser Phe Ser Gln Asn Pro Pro Val Leu Lys Arg His Gln Arg

1

5

10

10

CLAIMS

We claim:

1. A method of treating hemophilia in a human comprising:
 - a) providing at least one recombinant adeno-associated virion comprising nucleotide sequences encoding Factor VIII; and
 - b) administering said at least one recombinant adeno-associated virion into said human under conditions such that said Factor VIII nucleotide sequences are expressed at a level which provides a therapeutically effective amount in said human.
2. The method of Claim 1, wherein said Factor VIII nucleotide sequences are expressed in the liver of said human.
3. The method of Claim 1, wherein said recombinant adeno-associated virion encodes the light chain of said Factor VIII.
4. The method of Claim 1, wherein said recombinant adeno-associated virion encodes the heavy chain of said Factor VIII.
5. The method of Claim 1, wherein said adeno-associated virion is administered intravenously.
6. The method of Claim 1, wherein the nucleic acid encodes the amino acid sequence set forth in SEQ ID NO:15.

7. A method of treating a subject comprising:

a) providing:

- i) a subject suffering from a blood clotting disorder,
- ii) a first recombinant adeno-associated virion, comprising a nucleotide sequence encoding the light chain of Factor VIII, and
- iii) a second recombinant adeno-associated virion comprising a nucleotide sequence encoding the heavy chain of Factor VIII; and

b) administering said first and second recombinant adeno-associated virions into said subject under conditions such that said Factor VIII heavy and light chain nucleotide sequences are expressed at a level which provides a therapeutic effect in said subject.

8. The method of Claim 7, wherein said recombinant adeno-associated virion is administered intravenously.

9. The method of Claim 7, the nucleic acid encodes the amino acid sequence set forth in SEQ ID NO:15.

10. A method of treating hemophilia in a mammal comprising:
- (a) providing
- i) a recombinant adeno-associated virion, wherein said recombinant adeno-associated virion comprises a nucleic acid encoding Factor VIII operably linked to an expression control element, and
- 5 ii) a mammal; and
- (b) administering an amount of said recombinant adeno-associated virion to a mammal wherein said Factor VIII is expressed at levels having a therapeutic effect on said mammal.
- 10
11. The method of Claim 10, wherein said nucleic acid encodes the light and heavy chains of Factor VIII.
12. The method of Claim 10, wherein said mammal is a human.
- 15
13. The method of Claim 10, wherein said administering comprises introducing said recombinant adeno-associated virion to the liver of said mammal.
14. The method of Claim 10, wherein said recombinant adeno-associated virion is introduced intravenously.
- 20
15. The method of Claim 14, wherein the recombinant adeno-associated virion is introduced via the portal vein.
- 25
16. The method of Claim 10, wherein said recombinant adeno-associated virion is introduced arterially.

17. The method of Claim 16, wherein said recombinant adeno-associated virion is introduced via the hepatic artery.

5 18. The method of Claim 10, wherein said nucleic acid comprises SEQ ID NO: 13.

19. The method of Claim 10, wherein said nucleic acid comprises SEQ ID NO: 14.

10 20. The method of Claim 10, wherein said expression control element is tissue-specific.

21. The method of Claim 10, wherein the expression control element is liver-specific.

15 22. The method of Claim 10, said nucleic acid encodes the amino acid sequence set forth in SEQ ID NO:15.

20 23. A method of treating hemophilia A in a mammal comprising:
(a) providing:
i) a recombinant adeno-associated virion wherein said recombinant adeno-associated virion comprises a nucleic acid encoding a Factor VIII lacking a portion of the B-domain region and wherein said nucleic acid is operably linked to an expression control element, and
25 ii) a mammal suffering from hemophilia A; and

(b) administering an amount of said recombinant adeno-associated virion to a mammal wherein said Factor VIII light chain and heavy chain are expressed at levels having a therapeutic effect on said mammal.

5

24. The method of Claim 23, wherein said nucleic acid encodes the light and heavy chains of Factor VIII.

25. The method of Claim 23, wherein said mammal is a human.

10

26. The method of Claim 23, wherein said administering step comprises introducing the recombinant adeno-associated virion to the liver of said mammal.

15

27. The method of Claim 23, wherein said recombinant adeno-associated virion is introduced intravenously.

28. The method of Claim 27, wherein said recombinant adeno-associated virion is introduced via the portal vein.

20

29. The method of Claim 23, wherein said recombinant adeno-associated virion is introduced arterially.

30. The method of Claim 29, wherein said recombinant adeno-associated is introduced via the hepatic artery.

25

31. The method of Claim 23, wherein said nucleic acid comprises the Factor VIII sequence of SEQ ID NO: 13.

32. The method of Claim 23, wherein said nucleic acid comprises the Factor VIII sequence of SEQ ID NO: 14.

33. The method of Claim 23, wherein said expression control element is tissue-specific.

34. The method of Claim 23, wherein the expression control element is liver-specific.

35. The method of Claim 34, wherein said Factor VIII nucleotide sequences are expressed in the liver of said subject.

36. The method of Claim 23, wherein said nucleotide sequences are operably linked to tissue-specific expression control elements.

37. The method of Claim 23, wherein said nucleotide sequences are operably linked to liver-specific expression control elements.

38. The method of Claim 23, wherein said Factor VIII nucleotide sequences are operably linked to a tissue-specific expression control element.

39. The method of Claim 38, wherein said Factor VIII nucleotide sequences are operably linked to a liver-specific expression control element.

40. The method of Claim 23, said nucleic acid encodes the amino acid sequence set forth in SEQ ID NO:15.

ABSTRACT

The present invention provides improved viral vectors useful for the expression of genes at high levels in human cells. In particular, the present invention provides recombinant adeno-associated vectors (AAV) suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of full-length therapeutic levels of biologically active Factor VIII in the recipient individual *in vivo*. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using these constructs.